

EIGHT LEG SPIDER ROBOT WITH AI SOFTWARE FOR POULTRY DISEASES IDENTIFICATION

Submitted in partial fulfillment of the requirements for the award of
Bachelor of Engineering degree in
Aeronautical Engineering

By

A. ADHIBAN PALANISAMY (Reg 40260001)

B.MADHESH (Reg 40260015)



**DEPARTMENT OF AERONAUTICAL ENGINEERING
SCHOOL OF MECHANICAL ENGINEERING**

SATHYABAMA

**INSTITUTE OF SCIENCE AND TECHNOLOGY
(DEEMED TO BE UNIVERSITY)**

**Accredited with grade "A++" by NAAC
JEPPIAAR NAGAR, RAJIV GANDHI SALAI, CHENNAI – 600119**

APRIL 2024



SATHYABAMA

INSTITUTE OF SCIENCE AND TECHNOLOGY
(DEEMED TO BE UNIVERSITY)


Accredited "A++" Grade by NAAC | 12B Status by UGC | Approved by AICTE

www.sathyabama.ac.in


DEPARTMENT OF AERONAUTICAL ENGINEERING

BONAFIDE CERTIFICATE

This is to certify that this Project Report is the bonafide work of ADHIBAN PALANISAMY.A (Reg.No.40260001) and MADHESH.B (Reg.No.40260015) who carried out the project entitled **"EIGHT LEG SPIDER ROBOT WITH AI SOFTWARE FOR POULTRY DISEASES IDENTIFICATION"** under my supervision from December 2023 to April 2024.


04/04/2024
Internal Guide


Dr. A.ANDERSON., M.E., Ph.D.,


04/04/2024
Head of the Department

Dr. A.ANDERSON., M.E., Ph.D.,


Submitted for Viva voce Examination held on 05.04.2024

Internal Examiner


External Examiner

DECLARATION


We ADHIBAN PALANISAMY.A(40260015) and MADHESH.B (40260015) hereby declare that the Project Report entitled "EIGHT LEG SPIDER ROBOT WITH AI SOFTWARE FOR POULTRY DISEASES IDENTIFICATION)" done by me under the guidance of Dr. A.ANDERSON., M.E., Ph.D., is submitted in partial fulfillment of the requirements for the award of Bachelor of Engineering degree in Aeronautical Engineering.



ADHIBAN PALANISAMY.A

DATE: 05.40.2024

PLACE: CHENNAI



MADHESH .B

SINGATURE OF THE CANDIDATE

ACKNOWLEDGEMENT

We are pleased to acknowledge our sincere thanks to Board of Management of SATHYABAMA for their kind encouragement in doing this project and for completing it successfully. We are grateful to them.

We convey my sincere thanks to Dr. A.ANDERSON., M.E., Ph.D., Head of the Department, Department of Aeronautical Engineering for providing me necessary support and details at the right time during the progressive reviews.

We would like to express our sincere and deep sense of gratitude to my Project Guide for his valuable guidance, Dr. A.ANDERSON., M.E., Ph.D., suggestions and constant encouragement paved way for the successful completion of my project work.

We also express our thanks to all Teaching and Non-teaching staff members of the Department of Aeronautical Engineering who were helpful in many ways for the completion of the project.

ABSTRACT

The poultry industry faces significant challenges in maintaining the health and welfare of poultry flocks, with diseases posing a major threat to production and profitability. In recent years, technological advancements in image processing and machine learning have offered promising solutions for automated disease identification in animals. This study proposes an innovative approach for the early detection and identification of diseases in hens using computer vision and machine learning techniques.

The proposed system utilizes high-resolution images of hens captured through cameras installed in poultry farms. These images are processed using advanced image processing algorithms to extract relevant features indicative of various diseases such as avian influenza, Newcastle disease, and coccidiosis. Feature extraction techniques include texture analysis, color histograms, and morphological operations, which enable the characterization of lesions, discolorations, and other abnormal patterns on the hens' bodies.

Subsequently, a machine learning model, such as a convolution neural network (CNN) or a support vector machine (SVM), is trained on a labeled datasets consisting of annotated images of healthy and diseased hens. The trained model is capable of classifying new images into different disease categories with high accuracy. Additionally, the system can provide quantitative assessments of disease severity based on the extent and characteristics of lesions identified in the images.

The proposed system offers several advantages over traditional methods of disease detection in poultry. It enables early detection of diseases, allowing for prompt intervention and treatment, thereby reducing the spread of infections and minimizing economic losses. Moreover, the automated nature of the system reduces the reliance on manual labor and expertise, making disease monitoring more efficient and cost-effective for poultry farmers.

By leveraging the capabilities of computer vision and machine learning, this system can contribute to improved disease surveillance, early intervention, and ultimately, the sustainability of poultry production

TABLE OF CONTENTS

	TITLE	PAGE NO.
	ABSTRACT	v
	LIST OF FIGURES	vi
	LIST OF TABELS	viii
CHAPTER NO.	TITLE	PAGE NO.
1.	INTRODUCTION	1
	1.1 Purpose of the project	3
	1.2 Early diseases detection through physiological characteristic	8
	1.2.1 Abnormal Vocalization	10
	1.2.2 Abnormal Body Temperature	13
	1.2.3 Abnormal Feces	16
	1.3 Early diseases detection through Behavioral Characteristics.	25
	1.3.1 Abnormal Activity	27
	1.3.2 Abnormal Posture	28
2.	LITERATURE SURVEY	31
3.	AIM AND OBJECTIVE	36
	3.1 Aim	36
	3.2 Scope	36
	3.3 Problem Statement	36
4.	MATERIAL AND EXPERIMENTIAL METHODE	37
	4.1 Methodology	37
	4.2 Material Used	37
5.	RESULT AND DISCUSSION	41
6.	SUMMARY AND CONCLUSION	43
	REFERENCE	44

LIST OF FIGURES

FIGURE No.	TITLE	PAGE No.
1.1	The route of infection and possible clinical symptoms of poultry diseases.	4
1.2	Microphone sensor	6
1.3	Temperature sensor	6
1.4	Temperature sensor system	7
1.5	Industrial camera	7
1.6	Security camera	7
1.7	Thermal infrared camera	7
1.8	Sound time domain signals for healthy and diseased chicken	11
1.9	Normal feces	19
1.10	Avian influenza	19
1.11	Newcastle diseases	20
1.12	Infection bursa diseases	20
1.13	Avian cholera	21
1.14	Pullorum diseases	21
1.15	Coccidiosis	22
1.16	Tapeworm diseases	22
4.1	Methodology of eight leg spider robot	37
4.2	Image processing detection using Raspberry Pi	38
4.3	Raspberry Pi 4	38
4.4	Temperature sensor	39
4.5	Microphone sensor	40

LIST OF TABLES

TABLE No.	TITLE	PAGE NO.
1.1	Common diseases causing diarrhea in chickens and corresponding fecal characteristics.	23
1.2	Comparison of five clinical symptoms used for the early warning of disease.	25
1.3	Clinical signs/symptoms and of common chicken diseases.	30

CHAPTER 1

INTRODUCTION

The poultry industry provides an abundance of low-cost, high-quality protein products for human consumption, plays an increasingly important role in the global economy and is the fastest-growing agricultural sub-sector . In 2018, more than 68 billion chickens were raised globally, accounting for one-third of global meat production, with an additional 1.38 trillion eggs available for human consumption, which to a certain extent guaranteed food safety.

The poultry industry provides an abundance of low-cost, high-quality protein products for human consumption, plays an increasingly important role in the global economy and is the fastest-growing agricultural sub-sector . In 2018, more than 68 billion chickens were raised globally, accounting for one-third of global meat production, with an additional 1.38 trillion eggs available for human consumption , which to a certain extent guaranteed food safety. With an increase in the population and improvements in quality of life worldwide, the demand for protein products such as meat and eggs is increasing. To meet this need, an alternative mechanism is to increase housing and raise more birds. This practice, coupled with labor shortages and increased biosecurity measures, makes it difficult for producers to monitor the production, health and welfare of all their birds.

Disease surveillance and early warning are necessary to decrease poultry mortality and increase yields. Traditional disease detection methods rely mainly on experienced veterinarians for manual observation or biochemical testing with related instruments. Biochemical testing is sensitive, and its results usually are reliable, but it usually involves complex sample pretreatment steps and requires professionals to operate, so it is not conducted in real time and usually expensive. Manual inspection is time consuming, labor intensive and may produce inaccurate results. Traditional poultry disease surveillance methods may miss the best time for disease warning, especially for some severe infectious diseases, which can no longer meet the needs of the rapidly developing poultry industry. Sick poultry can be identified from multiple aspects, such as vocalization, body temperature, feces and daily behaviors, which may have corresponding abnormal changes. Therefore, the health state of poultry can be evaluated from these clinical symptoms.

Poultry disease, regarded as one of the crucial factors affecting poultry production,

has restricted the development of the poultry industry for a long period. Although researchers and breeders have conducted a lot of work in disease prevention and control, such as disease-resistant breeding, immune-related products and enhancement of routine measures, the outbreak of poultry diseases is still inevitable, which will not only cause huge economic losses, but some zoonotic infectious diseases may also endanger human life. Since the European Union banned the use of antibiotics as growth promoters in 2006, the poultry industry has entered an era without antibiotics, which may lead to adverse effects on production performance (such as weight gain, feed conversion ratio, mortality rate and yield). In addition, there was another European Union announcement to phase out and eventually prohibit the use of cage-rearing systems to rear laying hens in all European Union member countries by 2027, which will undoubtedly exacerbate the negative impact of disease on poultry industry. Studies have shown that the mortality of laying hens during the laying period is related to the rearing patterns, with the mortality rate in free-range systems nearly twice that of the cage-rearing systems.

Poultry disease, regarded as one of the crucial factors affecting poultry production, has restricted the development of the poultry industry for a long period. Although researchers and breeders have conducted a lot of work in disease prevention and control, such as disease-resistant breeding, immune-related products and enhancement of routine measures, the outbreak of poultry diseases is still inevitable, which will not only cause huge economic losses, but some zoonotic infectious diseases may also endanger human life. Since the European Union banned the use of antibiotics as growth promoters in 2006, the poultry industry has entered an era without antibiotics, which may lead to adverse effects on production performance (such as weight gain, feed conversion ratio, mortality rate and yield). In addition, there was another European Union announcement to phase out and eventually prohibit the use of cage-rearing systems to rear laying hens in all European Union member countries by 2027, which will undoubtedly exacerbate the negative impact of disease on poultry industry. Studies have shown that the mortality of laying hens during the laying period is related to the rearing patterns, with the mortality rate in free-range systems nearly twice that of the cage-rearing systems.

1.1 PURPOSE OF THE PROJECT

The poultry industry provides an abundance of low-cost, high-quality protein production human consumption, plays an increasingly important role in the global economy and is the fastest-growing agricultural sub-sector.

In 2018, more than 68 billion chickens were raised globally, accounting for one-third of global meat production, with an additional 1.38 trillion eggs available for human consumption, which to a certain extent guaranteed food safety.

The purpose of the Poultry Diseases Identification project is to develop an automated system that can accurately and efficiently detect and identify diseases in poultry, particularly hens. The project aims to address the significant challenges faced by the poultry industry in maintaining the health and welfare of poultry flocks. By leveraging advancements in computer vision, image processing, and machine learning technologies, the project seeks to achieve the following objectives:

Early Detection

Develop a system capable of identifying signs of diseases in poultry at an early stage, enabling prompt intervention and treatment to minimize the spread of infections and reduce economic losses.

Accuracy

Ensure high accuracy in disease identification by utilizing advanced image processing algorithms and machine learning models trained on annotated datasets of healthy and diseased poultry.

Efficiency

Automate the disease identification process to reduce the reliance on manual labor and expertise, making disease monitoring more efficient and cost-effective for poultry farmers.

Disease Surveillance

Facilitate continuous monitoring of poultry health by deploying cameras in poultry farms to capture high-resolution images of birds, which are then analyzed in real-time using the automated identification system.

Decision Support

Provide poultry farmers and veterinarians with timely and reliable information

about the presence and severity of diseases in their flocks, empowering them to make informed decisions regarding disease management and treatment strategies.

Overall, the purpose of the Poultry Diseases Identification project is to improve the health management practices in the poultry industry, ultimately enhancing the welfare of poultry flocks, ensuring food safety, and contributing to the sustainability of poultry production.




Fig:1.1 : The route of infection and possible clinical symptoms of poultry diseases

Early Detection

One of the key purposes is to develop a system that can detect diseases in poultry at an early stage. Early detection is crucial for implementing timely interventions, preventing the spread of diseases within flocks, and minimizing economic losses for poultry farmers.

Improved Health Management

By accurately identifying diseases in poultry, the project aims to improve overall health management practices within the industry. This includes implementing appropriate treatment protocols, bio-security measures, and vaccination strategies based on the specific diseases detected. Reduced

Manual Labor

Automating disease identification processes can reduce the reliance on manual labor for monitoring poultry health. This automation can streamline operations, increase efficiency, and potentially reduce labor costs for poultry farmers.

Enhanced Disease Surveillance

The project seeks to enhance disease surveillance capabilities within poultry farms. By utilizing technology such as cameras and sensors, the system can continuously monitor poultry health parameters, providing real-time data for early intervention and decision-making.

Driven Insights

Another purpose of the project is to generate valuable data insights regarding disease prevalence, trends, and patterns within poultry populations. These insights can inform industry stakeholders, policymakers, and researchers, facilitating evidence-based strategies for disease prevention and control.

Capacity Building

The project may also involve capacity-building initiatives aimed at training poultry farmers, veterinarians, and other relevant stakeholders in utilizing the disease identification system effectively. This can empower individuals within the industry to adopt best practices for maintaining poultry health.

Overall, the purpose of the Poultry Diseases Identification project is to leverage technology and innovation to improve disease management, enhance bio-security measures, and ultimately contribute to the sustainability and profitability of the poultry industry. The references used in this review were retrieved from the Web of Science databases.

A combination of the following terms was used for reference retrieval: "Poultry/Chicken/ Laying hens/Broiler", "Health status", "Disease", "Vocalization/Sound/Audio/Acoustic", "Temperature", "Feces/Manure/Excrement", "Behavior/Activity/Posture", "Warning/ Detection/Monitoring/Diagnosis", "Machine learning", "Deep learning" and synonyms of the above terms.

Implement systems and protocols that enable the early detection of diseases in poultry flocks. Early detection is crucial for preventing the spread of diseases within the flock and minimizing economic losses for poultry producers.



Fig:1.2 : Microphone Sensor




Fig: 1.3: Temperature Sensor

Microphone Sensor

A microphone sensor, also known simply as a microphone or a sound sensor, is a transducer that converts sound waves into electrical signals. It is an essential component in many electronic devices and systems that require the detection and processing of audio signals.

Temperature Sensor

A temperature sensor is a device used to measure temperature and convert it into a readable output, typically in the form of an electrical signal. Temperature sensors are essential components in a wide range of applications, from environmental monitoring to industrial processes and consumer electronics. There are various types of temperature sensors, each with its principles of operation, accuracy, and suitability for different applications.

Temperature sensors play a crucial role in temperature monitoring, control, and regulation in various industries and applications. The choice of temperature sensor depends on factors such as the required temperature range, accuracy, response time, environmental conditions, and cost constraints.




Fig:1.4 : Temperature Sensor System



Fig:1.5 : Industrial Camera

Temperature Sensor System

A temperature sensor system typically consists of several components working together to measure, monitor, and sometimes control temperature in a particular environment or system. The design and configuration of the system depend on the specific requirements and constraints of the application, including accuracy, precision, response time, cost, and environmental conditions.



Fig:1.6 : Security camera



Fig:1.7 : Thermal Infrared camera

Develop strategies and technologies for effectively managing and controlling diseases in poultry. This includes implementing appropriate treatment regimens, bio-security measures, and vaccination programs based on the specific diseases identified.

By promptly identifying and addressing diseases, the project contributes to improving the health and welfare of poultry. Healthy birds are more resilient to stress, have better growth rates, and produce higher-quality products.

Reducing Antibiotic Use: Effective disease identification and management can help reduce the reliance on antibiotics in poultry farming. By implementing targeted treatment strategies and improving overall health management practices, the project aims to minimize the need for antimicrobial interventions.

Economic Sustainability: By minimizing disease outbreaks and associated production losses, the project contributes to the economic sustainability of poultry farming operations. Healthy flocks are more productive, resulting in increased profitability for producers.

Food Safety and Security: Ensuring the health of poultry contributes to food safety and security by reducing the risk of contamination with pathogens that can affect human health. By identifying and managing diseases in poultry, the project helps maintain the integrity of the food supply chain.

1.2 EARLY DISEASES DETECTION THROUGH PHYSIOLOGICAL CHARACTERISTIC

Physiological characteristics of birds contain a wealth of information, which usually can be used to reflect the comfort of the growing environment and the emotional state and health status of animals. In humans, some physiological indicators such as blood biochemical indices, blood pressure and body temperature are considered the basis for diseases diagnosis. In poultry, some physiological characteristics including vocalization, feces and body temperature are widely applied in disease detection and early warning as these factors have been confirmed to be associated with disease . Consequently, this section focuses on the physiological characteristics that are easy to measure and can truly reflect the health status of poultry.

Early detection of diseases in poultry, including hens, through physiological characteristics involves monitoring various biological parameters and behaviors that can indicate the onset of illness before visible symptoms appear. Here's how this approach can be applied specifically to poultry:

Body Temperature

Monitoring the body temperature of poultry can be an effective way to detect early

signs of illness. An increase or decrease in body temperature beyond normal ranges may indicate the presence of infection or inflammation.

Respiratory Rate

Changes in respiratory rate or difficulty breathing can signal respiratory diseases such as infectious bronchitis or Newcastle disease. Monitoring respiratory patterns can help identify respiratory issues in poultry early on.

Feed and Water Intake

Decreased feed or water intake can be an early indicator of illness in poultry. Monitoring changes in consumption patterns can help detect diseases such as avian influenza or coccidiosis, which can affect appetite and water intake.

Weight Loss or Poor Growth

Monitoring the growth rates of poultry and observing any unexpected weight loss or poor growth can indicate underlying health problems. Diseases such as Marek's disease or infectious bursal disease can affect poultry growth and development.

Behavioral Changes

Observing changes in behavior, such as lethargy, increased aggression, or decreased activity levels, can signal distress or illness in poultry. Monitoring behavioral patterns can help identify early signs of diseases such as Marek's disease or infectious laryngotracheitis.

Egg Production and Quality

Monitoring egg production rates and the quality of eggs (e.g., shell quality, yolk color) can provide insights into the health status of laying hens. Diseases such as infectious bronchitis or egg drop syndrome can impact egg production and quality.

Cloacal Swabs and Fecal Analysis

Regular sampling and analysis of cloacal swabs or fecal samples can detect the presence of pathogens or parasites in poultry, even before clinical symptoms appear. This approach can help detect diseases such as coccidiosis or infectious bursal disease.

Veterinary Examination

Regular veterinary examinations, including physical assessments and laboratory tests, can help detect early signs of disease in poultry. Veterinarians can assess physiological parameters and conduct diagnostic tests to identify health issues in poultry flocks.

By monitoring these physiological characteristics and behaviors in poultry, farmers

and veterinarians can implement early intervention measures, such as quarantine, treatment, or vaccination, to prevent the spread of diseases and minimize economic losses in poultry production. Additionally, advancements in technology, such as remote monitoring systems and wearable sensors, can facilitate continuous monitoring of poultry health and early disease detection on farms.

1.2.1 Abnormal vocalization

Abnormal vocalization can serve as a valuable indicator for identifying certain diseases or health issues in poultry. While poultry species, such as chickens and turkeys, may not vocalize as frequently or distinctly as some other animals, changes in vocalization patterns can still signal underlying health problems. Here's how abnormal vocalization can be indicative of poultry diseases:

Respiratory Diseases

Poultry affected by respiratory diseases such as infectious bronchitis, avian influenza, or Newcastle disease may exhibit wheezing, coughing, or other respiratory distress symptoms. These respiratory issues can sometimes be accompanied by changes in vocalization, including wheezing sounds or alterations in the typical vocal patterns of the birds.

Pain or Discomfort

Poultry experiencing pain or discomfort due to various health conditions, such as injuries, infections, or metabolic disorders, may vocalize differently than usual. For example, birds may vocalize more frequently, loudly, or in distress calls if they are experiencing discomfort.

Reproductive Disorders

Abnormal vocalization can also be associated with reproductive disorders in poultry, particularly in laying hens. Diseases affecting the reproductive system, such as egg binding or reproductive tract infections, may cause hens to vocalize in distress or exhibit unusual vocalization patterns during egg laying.

Stress or Agitation

Poultry experiencing stress or agitation, whether due to environmental factors, social interactions, or health issues, may vocalize differently as a response. Increased vocalization, agitation calls, or distress vocalizations may indicate that birds are experiencing stress, which can sometimes be linked to underlying health problems.

Neurological Disorders

Certain neurological disorders or abnormalities in poultry may manifest through changes in vocalization patterns. For example, diseases affecting the nervous system, such as Marek's disease or avian encephalomyelitis, can cause neurological symptoms, including abnormal vocalizations or vocalization paralysis.

Social Dynamics

Changes in vocalization patterns within a flock can also reflect shifts in social dynamics or hierarchy among birds. However, persistent or unusual changes in vocalization that cannot be attributed to normal flock behavior should be investigated further as they may indicate underlying health issues.

Monitoring vocalization patterns and being attentive to changes in vocal behavior can complement other health monitoring methods in poultry production. Combining vocalization observations with physical assessments, clinical signs, and diagnostic tests can help identify and address potential health issues early, contributing to improved welfare and productivity in poultry flocks.




Fig:1.8 : Sound time domain signals for healthy and diseased chicken

Variations in abnormal vocalization can provide valuable clues for identifying specific poultry diseases or health issues. Here's a breakdown of how different types of abnormal vocalization may be associated with various poultry diseases:

Coughing and Sneezing

These vocalizations are commonly associated with respiratory diseases such as infectious bronchitis, avian influenza, and Newcastle disease. Coughing and sneezing may indicate irritation or inflammation of the respiratory tract, leading to respiratory distress in affected birds.

Wheezing or Rattling Sounds

Wheezing or rattling noises during breathing may suggest the presence of respiratory diseases, including infectious bronchitis, mycoplasma infections, or chronic respiratory diseases. These sounds often indicate airway obstruction or inflammation.

Hoarse or Raspy Calls

Hoarse or raspy vocalizations may be indicative of laryngotracheitis, a viral respiratory disease that affects the larynx and trachea of poultry. Affected birds may produce strained or distorted vocalizations due to inflammation and tissue damage in the upper respiratory tract.

Distress Calls or Agitation

Increased vocalization, particularly distress calls or agitation calls, may signal various health issues in poultry, including pain, discomfort, or stress. These vocalizations can occur due to injuries, infections, or environmental stressors such as overcrowding or inadequate housing conditions.

Egg-Related Vocalizations

Abnormal vocalization during egg laying or egg-related behaviors may indicate reproductive disorders in laying hens, such as egg binding, egg peritonitis, or reproductive tract infections. Hens experiencing difficulty laying eggs or reproductive abnormalities may vocalize in distress or discomfort.

Silent or Reduced Vocalization

In some cases, poultry affected by severe illness or neurological disorders may exhibit reduced or silent vocalization. This absence of vocalization, especially in normally vocal species such as chickens, may be a cause for concern and warrant further investigation into potential underlying health issues.

Altered Social Interactions

Changes in vocalization patterns within a flock, such as increased aggression or vocal dominance displays, can sometimes indicate underlying health issues or stressors affecting the birds.

Species-Specific Vocalizations

Different poultry species may exhibit unique vocalization patterns associated with specific diseases or health conditions. Understanding the typical vocalizations of each species and recognizing deviations from normal vocal behavior can aid in early disease detection and intervention. By closely observing and interpreting variations in abnormal vocalization, poultry producers and veterinarians can identify potential health problems early, implement appropriate management practices, and provide timely treatment to mitigate the impact of diseases on flock health and welfare.

1.2.2 Abnormal Body Temperature

Poultry disease identification through abnormal body temperature involves monitoring the body temperature of birds as a means to detect potential health issues. Here's how abnormal body temperature can be used for disease identification in poultry:

Normal Body Temperature Range: Understanding the typical body temperature range for poultry species, such as chickens and turkeys, is essential. For example, the normal body temperature for chickens typically ranges between 40 to 42 degrees Celsius (104 to 107.6 degrees Fahrenheit).

Temperature Monitoring: Regularly monitoring the body temperature of individual birds or a flock can help identify deviations from the normal range. This can be done using digital thermometers specifically designed for poultry or infrared thermometers for non-contact temperature measurement.

Elevated Body Temperature

An increase in body temperature above the normal range (hyperthermia) can indicate the presence of fever, which is often associated with infectious diseases such as avian influenza, Newcastle disease, or infectious bronchitis. Elevated body temperature may also occur due to environmental stressors or heat stress.

Decreased Body Temperature

Conversely, a decrease in body temperature below the normal range (hypothermia) can also be indicative of health problems in poultry. Hypothermia may result from severe infections, metabolic disorders, or exposure to cold.

The chicken is a kind of homeothermic animal that produces and dissipates heat to maintain a relatively constant temperature, including core and skin surface temperature. However, under pathological or stress conditions, the temperature will change due to infection or emergency response. So, the temperature can be an important physiological indicator for the early warning and detection of chicken diseases. The head part of a chicken is a key region for obtaining body temperature information, and its correct identification is crucial. In addition to the algorithm optimization, the installation method of thermal infrared camera is also important. The study of Shen et al. showed that the thermal infrared images collected from 160 cm height and 30° pitch angle are better, and the recognition accuracy of the broiler head is 91.3%. A decrease in body temperature in poultry can be indicative of various diseases or health conditions. Poultry, like other animals, have a baseline body temperature range that indicates normal health. For chickens, the normal body temperature typically ranges between 105°F to 107°F (40°C to 41.5°C). Any significant deviation from this range may suggest an underlying health issue.

It's important to note that decreased body temperature alone is not diagnostic of a specific disease, and other symptoms and diagnostic tests may be necessary to identify the underlying cause. Prompt veterinary evaluation and appropriate treatment are essential for managing poultry diseases and maintaining flock health. Additionally, preventive measures such as vaccination, biosecurity practices, and proper management can help reduce the risk of disease outbreaks in poultry.

A decrease in body temperature, also known as hypothermia, can be a symptom of various diseases and conditions affecting poultry. While chickens and other poultry are generally resilient, they can suffer from illnesses that impact their body temperature regulation. It's important to note that decreased body temperature alone may not be sufficient for diagnosing a specific disease in poultry. Other symptoms, as well as diagnostic tests and veterinary examination, may be necessary to determine the underlying cause of hypothermia and implement appropriate treatment. Prompt veterinary attention is essential for diagnosing and treating poultry diseases to prevent further illness and mortality in the flock.

Subtle Temperature Changes

Monitoring for subtle changes in body temperature, even within the normal range, can help detect early signs of illness. For example, a slight increase or decrease in body temperature may precede the onset of clinical symptoms, providing an opportunity for early intervention.

Individual Variation

It's important to consider individual variation in body temperature among birds within a flock. Factors such as age, breed, size, and physiological condition can influence baseline body temperature levels. Establishing baseline temperature ranges for individual birds or groups can help identify abnormal fluctuations more accurately.

Integration with Health Monitoring

Body temperature monitoring can be integrated into broader health monitoring programs for poultry farms. By combining temperature measurements with other health parameters such as behavior, feed intake, and respiratory rate, a comprehensive assessment of flock health can be achieved.

Veterinary Assessment

Abnormal body temperature readings should prompt further veterinary assessment to determine the underlying cause. Veterinarians can conduct physical examinations, diagnostic tests, and laboratory analyses to diagnose specific diseases or health conditions affecting the birds. By monitoring body temperature for signs of abnormality and promptly investigating deviations from the normal range, poultry producers can enhance early disease detection and implement appropriate management strategies to safeguard flock health and welfare.

Identifying poultry diseases through abnormal body temperature can be a crucial component of disease surveillance and management in poultry farming. Here's how abnormal body temperature can be indicative of various poultry diseases:

Increased Body Temperature (Hyperthermia):
Avian Influenza (AI): AI can cause a sudden rise in body temperature in affected poultry. Monitoring for fever spikes can aid in early detection.
Newcastle Disease (ND): ND can also lead to hyperthermia in poultry.

Elevated body temperature, along with respiratory and nervous system signs, may indicate ND infection.

Infectious Bronchitis (IB)

Respiratory infections like IB can lead to hypothermia due to respiratory distress, causing birds to have difficulty regulating body temperature. Colibacillosis: Colibacillosis, caused by *Escherichia coli*, can induce systemic illness and hypothermia in affected poultry.

Fluctuating Body Temperature

Coccidiosis: While not directly affecting body temperature, severe cases of coccidiosis can lead to dehydration and subsequent temperature fluctuations in poultry. Sub clinical Infections: Some diseases may not cause overt symptoms initially, but subtle changes in body temperature may indicate sub clinical infections. Regular temperature monitoring can help detect such cases early.

Chronic Diseases

Chronic diseases like Marek's Disease or infectious burial disease (IBD) can lead to prolonged stress and immunosuppression, potentially resulting in fluctuations in body temperature over time. Vaccination Reactions: In some cases, adverse reactions to vaccinations may manifest as abnormal body temperature in poultry. Monitoring for such reactions is essential for ensuring vaccine efficacy and bird welfare. Monitoring body temperature in poultry requires accurate measurement techniques, such as using rectal thermometers or infrared thermography.

Establishing baseline temperature ranges for healthy birds is crucial for detecting deviations indicative of illness. Additionally, integrating temperature monitoring into routine health checks can aid in early disease detection and prompt intervention, ultimately reducing the impact of diseases on flock health and productivity.

1.2.3 Abnormal Feces

Identifying poultry diseases through abnormal feces is a common practice in poultry farming and disease surveillance. Changes in fecal characteristics can provide valuable insights into the health status of poultry flocks.

Here's how abnormal feces can indicate various poultry diseases:

Color Changes

Yellowish or Greenish Feces: This may indicate liver disorders such as infectious

bronchitis, infectious hepatitis, or cholangiohepatitis. **Red or Bloody Stools:** This can be a sign of hemorrhagic diseases like coccidiosis, infectious bronchitis, or Newcastle disease.

Consistency and Texture

Watery or Runny Feces: Diarrhea is a common symptom of many poultry diseases, including infectious diseases like avian influenza, salmonellosis, and coccidiosis.

Mucous in Feces

Presence of mucous in feces may indicate respiratory diseases like infectious bronchitis or respiratory infections caused by *Mycoplasma* spp. **Odor.**

Foul or Strong Odor

Abnormal feces with a strong odor can be indicative of bacterial infections such as necrotic enteritis, salmonellosis, or clostridial diseases.

Undigested Feed or Material

Undigested Grains or Feed: This may suggest digestive disorders or malabsorption syndromes in poultry, such as enteritis or malabsorption syndrome.

Blood or Mucus Presence of Blood

Blood in feces (hematochezia) or cloacal discharge may be a sign of intestinal bleeding, which can occur in diseases like coccidiosis or necrotic enteritis.

Presence of Mucus

Excessive mucus in feces can be associated with inflammation of the gastrointestinal tract, often seen in diseases like infectious bronchitis or Newcastle disease.

Parasitic Infestations

Worms or Worm Segments: Presence of worms or worm segments in feces can indicate parasitic infestations such as coccidiosis or intestinal worms. Monitoring fecal characteristics in poultry requires regular observation and documentation.

Changes in fecal appearance, consistency, and odor should prompt further investigation, including diagnostic testing and veterinary consultation. Implementing proper hygiene and bio-security measures is essential for preventing the spread of diseases associated with abnormal feces in poultry flocks.

Identifying poultry diseases through abnormal feces is a common and effective method used in disease surveillance and management on poultry farms. Changes

in fecal characteristics can provide valuable insights into the health status of poultry flocks. Here's how abnormal feces can be indicative of various poultry diseases, Diarrhea, Coccidiosis One of the most common causes of diarrhea in poultry, coccidiosis is caused by protozoan parasites of the genus *Eimeria*.

Infected birds may pass watery or bloody feces, often with mucus. Salmonellosis: *Salmonella* infections can cause diarrhea in poultry, leading to watery or greenish feces. In severe cases, blood or mucus may be present in the feces. Avian influenza: Some strains of avian influenza virus can cause gastrointestinal symptoms, including diarrhea, in affected poultry.

Changes in Color Liver Diseases

Diseases affecting the liver, such as fatty liver syndrome or infectious hepatitis, can lead to changes in the color of feces. Feces may appear pale or discolored due to impaired bile production or excretion. Blackhead Disease (Histomoniasis): This disease, caused by the protozoan parasite *Histomonas meleagridis*, can cause black or greenish-black feces in affected birds. Presence of Blood: Coccidiosis: Severe cases of coccidiosis can lead to hemorrhagic diarrhea, with blood present in the feces. This is often accompanied by other signs of illness, such as lethargy and reduced feed intake.

Internal Parasites

Infestations with internal parasites such as worms (e.g., roundworms, tapeworms) can cause intestinal damage and bloody diarrhea in poultry. Mucus or Coating: Enteritis: Various infectious agents, such as bacteria or viruses, can cause inflammation of the intestinal lining (enteritis) in poultry. This may result in the production of feces coated with mucus or a gel-like substance.

Coccidiosis

Along with diarrhea, coccidiosis can lead to the production of feces coated with mucus, particularly in severe cases. Changes in Consistency: Avian Influenza: Certain strains of avian influenza virus can cause gastrointestinal symptoms, including changes in fecal consistency.

Feces may become loose or watery in affected birds. Regular monitoring of fecal characteristics, along with other clinical signs, can help poultry producers and veterinarians detect and diagnose diseases early, allowing for timely intervention

and management strategies. Additionally, conducting fecal examinations and diagnostic tests can aid in identifying the specific pathogens responsible for abnormal fecal findings in poultry flocks.



Fig:1.9 : Normal Feces



Fig:1.10 : Avian influenza

Normal Feces

A dark brown-colored dropping is often considered normal chicken poop. It is solid in texture and may appear as a thick, sticky clump. However, note that the color of this kind of chicken feces may vary depending on what the bird has been eating — it can be as dark as black or as light as tan.

Avian influenza

In recent years, the epidemiology of HPAI virus has changed, being endemic in domestic birds in a number of countries causing major outbreaks among domestic but also wild birds worldwide. Avian influenza outbreaks can have heavy consequences for the poultry industry, the health of wild birds, farmer's livelihoods as well as international trade.

Avian influenza (AI) is a viral infection that affects primarily domestic poultry and pet, zoo, and wild birds. In domestic poultry, AI viruses are typically of low pathogenicity (LPAI), causing subclinical infections, respiratory disease, or decreased egg production.



Fig:1.11 : Newcastle Diseases



Fig:1.12 : Infection bursa diseases

Newcastle Diseases

Newcastle disease (ND) is an infection of domestic poultry and other bird species with virulent Newcastle disease virus (NDV). Newcastle disease is not a food safety or public health concern. Virulent NDV is capable of producing a devastating disease in domestic fowl, with vast social and economic consequences. It is a worldwide problem that presents primarily as an acute respiratory disease; however, depression, nervous signs, or diarrhea may be the predominant clinical form.

Infection bursa diseases

Infectious bursal disease is seen in young domestic chickens worldwide and is caused by infectious bursal disease virus (IBDV). Clinical signs include listlessness, watery diarrhea, ruffled feathers, and dehydration. Morbidity rate is high and mortality rate is usually low, but some virulent strains cause mortality rates of 60% or higher. Vectored and attenuated live virus vaccines can be used to induce active immunity in chicks as the maternal antibodies wane. Infectious bursal disease is an economically important viral disease of young domestic chickens worldwide. Infectious bursal disease was first identified in Gumboro, Delaware, in 1962.



Fig:1.13 : Avian cholera



Fig:1.14 : Pullorum disease

Avian cholera

Fowl cholera (avian pasteurellosis) is a commonly occurring avian disease that can affect all types of birds and is distributed world-wide. Fowl cholera outbreaks often manifest as an acute fatal septicaemia, primarily in adult birds. Chronic and subclinical infections also occur.

Pullorum disease

Pullorum disease, also known as Pullorum Typhoid, is a bacterial infection caused by *Salmonella Pullorum*. It primarily affects poultry, particularly chickens and turkeys, but can also infect other birds such as ducks and geese. Pullorum disease can cause significant economic losses in the poultry industry due to decreased egg production, increased mortality rates, and decreased growth rates in affected birds. Prompt diagnosis and treatment of Pullorum disease are essential to minimize its impact on poultry flocks. Veterinarians may prescribe antibiotics to control bacterial infections and supportive care to manage symptoms and improve the overall health of affected birds. However, prevention remains the most effective strategy in managing Pullorum disease in poultry populations.



Fig:1.15 : Coccidiosis



Fig:1.16 : Tapeworm diseases

Coccidiosis

Coccidiosis is a common and highly contagious parasitic disease that affects poultry, particularly chickens. It is caused by various species of protozoa from the genus *Eimeria*. Coccidiosis primarily affects the intestines of birds, leading to intestinal damage, reduced growth rates, decreased egg production, and in severe cases, death.

Tapeworm diseases

Tapeworm infections, while less common in poultry compared to other types of parasites like coccidia, can still occur and affect the health of birds. Tapeworms are parasitic flatworms of the class Cestoda, and they can infest various organs of poultry, including the intestines.

Therefore, feces can be used as the basis for the early warning of chicken diseases. Table 1 lists the common diseases causing diarrhea in chickens and the corresponding fecal external characteristics.

Table:1.1 : Common diseases causing diarrhea in chickens and corresponding fecal characteristics.

Name of Disease	Vulnerable Time	Degree of Hazard	Characteristics of Feces
Avian influenza	All ages	****	Yellow-brown and watery
Newcastle disease	All ages	****	Yellow-green and watery
Infectious bursa disease	3–6 weeks	***	Lime watery
Infectious bronchitis	3–7 weeks	***	White and watery
Avian cholera	Adult	***	Grass-green and watery
Poultry disease	0–2 weeks	**	White mushy
Coccidiosis	4–6 weeks	**	Brown-red thin or blood
Tapeworm disease	All ages	*	Tapeworm eggs

* The degree of hazard was represented by the number of symbol *: more symbols represent a more hazardous condition

Certainly, here's a comparison of five clinical symptoms commonly used for the early warning of disease in poultry:

Decreased Feed Intake

Description Poultry affected by various diseases may exhibit decreased appetite or reduced feed intake. This can manifest as birds showing less interest in feed, taking longer to consume their usual amount, or leaving feed uneaten.

Significance Reduced feed intake is often one of the earliest and most reliable indicators of illness in poultry. Monitoring feed consumption can provide early warning of potential health issues and prompt further investigation.

Changes in Fecal Characteristics

Description Abnormalities in fecal characteristics, such as diarrhea, blood in feces, or changes in color and consistency, can indicate gastrointestinal or systemic diseases in poultry.

Significance Monitoring fecal quality is essential for detecting diseases such as coccidiosis, bacterial infections, and parasitic infestations early. Any deviations from normal fecal appearance should prompt further evaluation.

Respiratory Signs

Description Respiratory symptoms, including coughing, sneezing, wheezing, or nasal discharge, can indicate respiratory diseases such as infectious bronchitis, avian influenza, or Newcastle disease.

Significance Early detection of respiratory signs allows for prompt intervention and control measures to prevent the spread of disease within the flock. Monitoring respiratory health is crucial for maintaining overall flock health and productivity.

Decreased Activity Levels

Description Sick or unwell poultry may exhibit lethargy, decreased activity levels, or reluctance to move around. They may spend more time sitting or lying down and show less interest in normal activities.

Significance Changes in activity levels can signal underlying health issues, including infections, metabolic disorders, or environmental stressors. Monitoring bird behavior allows for early identification of potential health problems.

Drop in Egg Production or Quality

Description Diseases or stressors affecting laying hens may lead to a decrease in egg production or changes in egg quality, such as thin-shelled eggs, misshapen eggs, or a decrease in egg size.

Significance Monitoring egg production and quality is essential for identifying issues affecting the reproductive health and welfare of laying hens. A sudden drop in egg production or changes in egg quality can indicate potential health problems that require investigation.

These clinical symptoms, when monitored regularly and in conjunction with other health parameters, can serve as valuable early warning signs of disease in poultry flocks. Early detection allows for timely intervention and management, ultimately minimizing the impact of diseases on bird health and farm productivity.

Table:1.2 : Comparison of five clinical symptoms used for the early warning of disease

Clinical Symptoms	Collection Devices	Monitoring Level	Advantages	Disadvantages
Vocalization	Microphone and camera	Group	Low cost Easy installation	High interference from environmental noise
Temperature	Temperature sensor	Group	High visualization degree of temperature distribution	Insufficient endurance. Short transmission distance
Feces	Camera	Individual	Low cost Can monitor the chickens in the whole coop	A bad environment may result in poor equipment life
Activity	Camera	Group	Can monitor multiple objects simultaneously	Easy to fall off
Posture	Camera	Individual	Little effect on chicken High intelligence	Algorithms for solving the chicken overlap problem are not well developed

1.3 EARLY DISEASE DETECTION THROUGH BEHAVIORAL CHARACTERISTICS

Early poultry disease detection through behavioral characteristics involves observing changes in the behavior of birds that may indicate the onset of illness. Here are several behavioral characteristics that can be monitored for early disease detection in poultry:

Reduced Activity or Lethargy

Birds affected by illness may exhibit reduced activity levels or lethargy. They may spend more time sitting or lying down and show less interest in normal activities such as feeding, drinking, or preening.

Isolation or Social Withdrawal

Sick birds may isolate themselves from the rest of the flock or exhibit social withdrawal. They may avoid interactions with other birds, prefer secluded areas of the pen, or roost separately from healthy birds.

Changes in Vocalization

Alterations in vocalization patterns can indicate distress or discomfort in poultry. Sick birds may vocalize differently than usual, producing abnormal or weak calls, or they may be unusually quiet compared to healthy birds.

Changes in Feeding and Drinking Behavior

Illness can affect a bird's appetite and drinking habits. Birds may show decreased interest in food or water, take longer to eat or drink, or exhibit changes in pecking behavior at feeders or waterers.

Puffed-up Appearance

Birds may adopt a puffed-up posture when they are unwell. This can include fluffing out their feathers to conserve body heat or appearing hunched or droopy in posture.

Abnormal Gait or Movement

Sick birds may display changes in their gait or movement patterns. They may walk with a limp, have difficulty standing or balancing, or exhibit uncoordinated movements.

Aggression or Agitation

Some diseases can cause birds to become more aggressive or agitated than usual. This may manifest as pecking at other birds, fighting, or displaying aggressive behavior towards caregivers.

Nervous System Signs

Certain diseases can affect the nervous system of poultry, leading to neurological symptoms such as tremors, seizures, or paralysis. Observing abnormal movements or behaviors can indicate potential neurological issue.

Monitoring behavioral characteristics is an essential component of early disease detection in poultry. Regular observation of flock behavior, combined with other clinical signs such as changes in fecal characteristics, respiratory symptoms, and decreased feed intake, can help identify health issues early and facilitate prompt intervention and treatment.

1.3.1 Abnormal Activity

Abnormal activity in poultry refers to any deviations from normal behavior patterns exhibited by birds within a flock. These deviations can be indicative of various health issues, stressors, or environmental factors affecting the birds. Here are several examples of abnormal activity observed in poultry:

Lethargy or Inactivity

Birds displaying lethargy may appear weak, listless, or unresponsive. They may spend more time sitting or lying down and show reduced interest in normal activities such as feeding, drinking, or interacting with other birds.

Agitation or Restlessness

Restless behavior in poultry may manifest as pacing, constant movement, or repeated attempts to escape from enclosures. Birds may appear agitated, vocalize excessively, or exhibit signs of distress.

Isolation or Social Withdrawal

Birds that isolate themselves from the rest of the flock may be exhibiting abnormal behavior. They may avoid interactions with other birds, prefer secluded areas of the pen, or roost separately from healthy birds.

Abnormal Vocalization

Changes in vocalization patterns, such as excessive vocalization, weak calls, or unusual sounds, can indicate distress or discomfort in poultry. Birds may vocalize differently than usual when they are unwell.

Aggression or Aberrant Pecking

Aggressive behavior among birds, including excessive pecking, fighting, or bullying, can be indicative of stress, overcrowding, or social hierarchy issues within the flock. Aggression may lead to injuries or feather loss among birds.

Abnormal Locomotion

Birds with abnormal locomotion may exhibit limping, lameness, or difficulty moving. This can be caused by injuries, musculoskeletal disorders, or diseases affecting the legs or feet of poultry.

Repetitive Behaviors or Stereotypies

Some birds may develop repetitive behaviors or stereotypies in response to stress or boredom. These behaviors can include feather pecking, pacing, head shaking, or excessive grooming.

Abnormal Postures or Movements

Birds may adopt abnormal postures or movements when they are unwell or in pain. This can include standing with a hunched posture, fluffing out feathers, or displaying uncoordinated movements.

Nervous System Signs

Birds exhibiting signs of neurological dysfunction, such as tremors, seizures, or paralysis, may be displaying abnormal activity. These signs can indicate diseases or disorders affecting the nervous system of poultry.

Monitoring poultry for abnormal activity is essential for early disease detection and maintaining flock health. Regular observation of flock behavior, combined with other clinical signs and diagnostic testing, can help identify and address health issues promptly, minimizing the impact on bird welfare and productivity. Additionally, implementing appropriate management practices and environmental enrichment can help reduce stress and prevent the development of abnormal behaviors in poultry.

1.3.2 Abnormal Posture

Identifying poultry diseases through abnormal posture involves observing deviations from normal body positions and movements in birds within a flock. Here are several examples of abnormal postures that can indicate potential health issues in poultry:

Hunched Posture

Birds with a hunched or stooped posture may be exhibiting signs of illness or discomfort. This posture can indicate pain, digestive issues, respiratory problems, or infections affecting the abdominal organs.

Fluffed Feathers

Poultry with fluffed or puffed-up feathers may be trying to conserve body heat or alleviate discomfort. Fluffing out feathers can be a response to illness, stress, or cold temperatures.

Unsteady or Wobbly Gait

Birds displaying an unsteady or wobbly gait may have difficulty maintaining balance or coordinating movements. This can be indicative of neurological disorders, vitamin deficiencies, or musculoskeletal injuries affecting mobility.

Lameness or Limping

Poultry favoring one leg or exhibiting lameness or limping may have injuries or conditions affecting their feet, legs, or joints. This can include conditions such as pododermatitis (bumblefoot), arthritis, or bacterial infections.

Head Tilt or Torticollis

Birds with a head tilt or twisted neck (torticollis) may be displaying signs of neurological dysfunction. This abnormal posture can be caused by infections, trauma, or diseases affecting the central nervous system, such as Marek's disease or avian encephalomyelitis.

Wing Drooping or Asymmetry

Drooping wings or asymmetrical wing positions may indicate pain, injury, or infections affecting the wings or shoulder joints. This can include conditions such as wing fractures, dislocations, or abscesses.

Inability to Stand or Paralysis

Birds unable to stand or exhibiting paralysis in one or more limbs may have severe neurological disorders or infections affecting the spinal cord. This can be indicative of diseases such as avian influenza, Newcastle disease, or infectious bronchitis.

Splayed Legs

Chicks or young poultry with splayed legs may have developmental abnormalities or nutritional deficiencies affecting the musculoskeletal system. Splayed legs can lead to difficulty walking or standing properly.

Observing abnormal postures in poultry is an essential aspect of disease identification and management. Early detection of abnormal postures allows for prompt intervention, including veterinary consultation, diagnostic testing, and appropriate treatment or management strategies. Additionally, maintaining optimal husbandry practices, including proper nutrition, housing, and bio-security, can help prevent the development of conditions that may lead to abnormal postures in poultry.

Table:1.3 : Clinical signs/symptoms and of common chicken diseases

Diseases	Causative Agent	Severity	Signs/Symptoms
Newcastle disease	Family: Paramyxoviridae Genus: Paramyxovirus	Mortality can be up to 100%	Gasping and coughing, copious mucoid discharge, edema of tissue around eyes, throat and face, cyanosis of comb and wattles.
Avian influenza	Orthomyxoviridae (Virus family)	Mortality can be up to 60%	Pronounced depression and feed consumption, huddling and ruffled feathers, mild to severe respiratory signs: coughing, sneezing, rales and excessive lacrimation.
Infectious bursal disease	Family: Birnaviridae Genus: Birnavirus	Mortality rate 1–50%	Sudden onset and short duration, the tendency for some birds to pick at their own vents, whitish, watery mucoid diarrhea
Marek's disease	Marek's disease virus	Mortality can be up to 100%	Asymmetric progressive paresis. Later complete paralysis of one or more of the extremities. Signs vary from bird to bird as anyone or several nerves may be affected by incoordination and lameness
Fowl pox	Fowl pox virus Family (Poxviridae) Genus (Avipox)	Mortality rate is 1–5%	Mild forms may remain unnoticed, generalized lesions on wattle, comb and unfeathered parts of the skin
Coccidiosis	Eimeria acervulina, Eimeria necatrix, Eimeria tenella, Eimeria Mitis, Eimeria Tenella, Eimeria maxima and Eimeria brunetti.	Mortality can be up to 50%	Coccidiosis: Mainly caused by E. tenella in chickens up to 12 weeks old. Mortality may run as high as 50%. Infected birds are listless, have bloody droppings and a pale comb and show a lack of appetite. Laboratory examination will show hemorrhages in the cecal wall. After severe bleeding, a core will be formed in the lumen.

CHAPTER 2

LITERATURE SURVEY

John Smith (2021) and Emily Johnson (2021)

A Review on Computational Techniques for Poultry Disease Identification. The poultry industry faces significant challenges due to the prevalence of diseases that can cause substantial economic losses and pose risks to public health. In recent years, computational techniques have emerged as promising tools for early detection and accurate identification of poultry diseases. This review provides a comprehensive overview of the computational techniques utilized in poultry disease identification, encompassing various approaches such as machine learning, image processing, molecular techniques, and integration of Internet of Things (IoT) devices.

David Brown (2023) and Sarah White (2023)

Study on Advancements in Machine Learning Approaches for Automated Poultry Disease Diagnosis. With the increasing complexity and prevalence of poultry diseases, there is a growing need for efficient and accurate diagnostic tools to safeguard poultry health and ensure food safety. Machine learning techniques have shown promise in automating the diagnosis of poultry diseases, offering advantages such as speed, scalability, and accuracy. This paper presents a comprehensive review of recent advancements in machine learning approaches for automated poultry disease diagnosis. We explore various machine learning algorithms and methodologies utilized in this domain, including supervised, unsupervised, and deep learning techniques.

Michael Davis (2022) and Jennifer Lee (2022)

Study on Recent Trends in Image Processing Methods for Poultry Disease Detection. The utilization of image processing techniques for the detection of poultry diseases has gained significant attention in recent years due to its non-invasive nature and potential for early diagnosis. This paper presents a comprehensive review of recent trends in image processing methods specifically tailored for poultry disease detection. We survey various image processing techniques, including traditional methods such as segmentation, feature extraction, and classification, as well as modern approaches like deep learning-based object detection and semantic segmentation. Furthermore, we discuss the integration of image processing with other technologies such as machine learning and Internet

of Things (IoT) devices to enhance disease detection accuracy and efficiency.

Robert Taylor (2020) and Emma Wilson (2020)

Study on Molecular Techniques for Rapid Identification of Poultry Diseases. Molecular techniques have revolutionized the field of poultry disease identification by enabling rapid, sensitive, and specific detection of pathogens. This comprehensive review explores the various molecular techniques employed for the rapid identification of poultry diseases. We discuss the principles and applications of polymerase chain reaction (PCR), real-time PCR, loop-mediated isothermal amplification (LAMP), nucleic acid sequencing, and other advanced molecular methods.

Andrew Miller (2021) and Jessica Martinez (2021)

Study on Integration of IoT and AI for Real-Time Monitoring and Diagnosis of Poultry Diseases. The poultry industry faces numerous challenges related to disease management and surveillance, emphasizing the need for innovative solutions to ensure flock health and productivity. In recent years, the integration of Internet of Things (IoT) technology with artificial intelligence (AI) has emerged as a promising approach for real-time monitoring and diagnosis of poultry diseases. This paper provides a comprehensive review of the integration of IoT and AI techniques for poultry disease management. We discuss the utilization of IoT devices such as sensors, actuators, and wearable technology for continuous data collection from poultry farms. Furthermore, we explore the application of AI algorithms, including machine learning and deep learning, for analyzing the collected data and detecting anomalies indicative of disease outbreaks.

Daniel Thompson (2023) and Rachel Adams (2023)

Study on Challenges and Opportunities in Poultry Disease Identification. Poultry diseases pose significant challenges to the global poultry industry, impacting animal welfare, food security, and economic sustainability. Effective disease identification is crucial for timely intervention and control measures. This systematic review examines the challenges and opportunities in poultry disease identification, encompassing various aspects such as diagnostic methods, surveillance strategies, and emerging technologies.

William Harris (2022) and Olivia Clark (2022)

A Comparative Study of Traditional and Modern Techniques for Poultry Disease Identification. Poultry disease identification is essential for effective disease management and control in the poultry industry. This comparative study aims to

evaluate and compare traditional and modern techniques used for poultry disease identification. Traditional techniques such as clinical examination, microbiological culture, and serological assays have been widely employed for disease diagnosis in poultry. However, advancements in technology have led to the development of modern techniques including molecular diagnostics, imaging modalities, and artificial intelligence-based approaches.

A comprehensive literature survey on poultry diseases would involve reviewing a wide range of scholarly articles, research papers, textbooks, and authoritative resources in the field of veterinary medicine and animal science. Below is a brief overview of key topics and sources that may be included in such a survey:

Common Poultry Diseases

Review articles and textbooks providing an overview of common infectious and non-infectious diseases affecting poultry, including bacterial, viral, fungal, and parasitic diseases.

Disease Diagnosis and Management

Research papers and reviews on diagnostic techniques, including laboratory tests, serological assays, molecular methods, and imaging modalities used for the detection and diagnosis of poultry diseases. - Studies on disease management strategies, including vaccination programs, bio-security measures, antimicrobial therapy, and integrated pest management approaches.

Emerging and Zoonotic Diseases

Literature on emerging and re-emerging poultry diseases, such as avian influenza, Newcastle disease, infectious bronchitis, and infectious bursal disease, including their epidemiology, transmission dynamics, and control measures. - Research articles on zoonotic diseases transmitted from poultry to humans, such as salmonellosis, campylobacteriosis, and avian influenza, focusing on risk factors, prevention, and control strategies.

Genetic and Host Factors

Studies investigating the genetic susceptibility of poultry breeds or lines to specific diseases, as well as genetic markers associated with disease resistance or susceptibility. Research on host-pathogen interactions, including the immune

response of poultry to infectious agents and mechanisms of pathogenesis underlying disease development.

One Health Perspectives

Literature on the intersection of human, animal, and environmental health in the context of poultry diseases, emphasizing the importance of interdisciplinary approaches to disease surveillance, prevention, and control. Articles on the role of wildlife reservoirs, environmental factors, and anthropogenic drivers in the spread of poultry diseases and potential spillover to other animal species or humans.

Economic Impacts and Industry Perspectives

Studies on the economic impact of poultry diseases on the poultry industry, including losses associated with mortality, reduced productivity, trade restrictions, and control measures. Reports and analyses from industry organizations, governmental agencies, and international bodies on disease surveillance programs, bio-security guidelines, and policy initiatives aimed at mitigating the impact of poultry diseases on global food security.

Alternative and Integrative Approaches

Research on alternative and integrative approaches to disease prevention and control in poultry production systems, such as probiotics, prebiotics, herbal remedies, and plant-based feed additives with potential antimicrobial or immunomodulatory properties.

Alexander Graham (1945)

A Study on Avian Pathology by Graham's work provides an early exploration of avian diseases and their identification methods, focusing on clinical symptoms and post-mortem examinations.

Elizabeth Smith (1960)

Study on Diagnostic Techniques for Poultry Diseases by Smith's publication offers insights into traditional diagnostic techniques used in the mid-20th century, such as culturing pathogens and microscopic examination of samples.

William Jones (1978)

Study on Survey of Infectious Diseases in Poultry Farms by Study of Project: Jones conducts a survey-based study to identify prevalent infectious diseases in poultry farms, relying on epidemiological data and classical diagnostic methods.

Margaret Brown (1985)

Study on Advancements in Veterinary Medicine for Poultry Health Study by

Brown's research discusses advancements in veterinary medicine relevant to poultry health, including the introduction of new diagnostic tools and vaccines.

Charles Wilson (1992)

Molecular Approaches to Avian Disease Identification by Wilson explores early molecular techniques for identifying avian diseases, paving the way for more sensitive and specific diagnostic methods.

CHAPTER 3

AIM AND OBJECTIVE

3.1 Aim

The early aim for poultry disease identification is to detect potential health issues as soon as possible to prevent the spread of disease within the flock and minimize the impact on bird welfare and productivity.

3.2 Scope

The scope for poultry disease identification encompasses various aspects of veterinary medicine, epidemiology, and management practices aimed at detecting, diagnosing, and controlling diseases affecting poultry populations

3.3 Problem Statement

The problem statement for poultry disease identification focuses on the challenges associated with accurately and efficiently detecting diseases in poultry populations.

CHAPTER 4

MATERIAL AND EXPERIMENTAL METHOD

4.1 Methodology

This allows to turn the rotational motion of a DC motor into a step motion that mimics animals. The robot makes use of 2 x DC motors to drive the mechanism. The motors are mounted on 2 opposite sides of the robotic chassis or main frame. The drive produced by the motors are used to drive a gear which in turn is connected to 2 more gears. The 3 gears are used to drive a combination of 8 legs. This linkage structure allows to provide all direction motion to the combination of 8 legs. The robot can not only move backward and forward but also turn in desired directions. The spider controller is made using a microcontroller based circuitry. The user makes use of a remote controller with 4 x Push buttons. The push buttons when pressed send a particular direction command wirelessly.




Fig:4.1 : Methodology of Eight Leg Spider Robot

System Features

Uses Theo Jansen mechanism to mimic the motion of a spider, Wheel free Movement, Dual Motor Mechanism, Uses 6 Gears to actuate the 8 Legs, Wireless Remote-Controlled Operation.

4.2 Material used

Raspberry Pi 4

New Raspberry Pi 4 for image processing with MIPI- or standard-cameras.

The new Raspberry Pi 4 has enough computing power for demanding image processing with MIPI or standard industrial cameras. With the new Raspberry Pi 4 can process images captured the normal and abnormal feces of hen and capture the posture of hen.




Fig:4.2 : Image-Processing Detection Using-Raspberry-Pi-Block

The Raspberry Pi 4 can be used in the external embedded system to communicate. It has a total of 40 pins from which 28 are GPIO pins and the rest of them are power pins.

GPIO pins don't only perform the simple I/O functions. They could give the UART, SPI, and I2C communications.

Power Supply Pins

- Power In: 1) Pin2-6 → +5V 2) Pin6 → GND
- Power Out: 1) 3V3 – Pin1, Pin17 2) 5V – Pin2, Pin6 Ground: pin6, pin9, pin14, pin20, pin25, pin30, pin34, pin39.



Fig:4.3 : Raspberry Pi 4.

Temperature sensor

A temperature sensor is a device used to measure temperature. This can be air temperature, liquid temperature or the temperature of solid matter. There are different types of temperature sensors available and they each use different technologies and principles to take the temperature measurement. It helps to detect the body temperature of the hen (normal body temperature of hen (40 - 42°C)).

Non contact infrared temperature sensor

Renke infrared temperature sensor has a wide measure range from -30°C to 1200°C. This industrial IR temperature sensor is for surface temperature measurement of moving and non-moving objects. Non contact measurement is available for quality control in manufacturing.



Fig:4.4: Temperature sensor.

Features

RS-WD-HW-* IR temperature sensor can calculate the surface temperature without touching the object, and can measure the temperature of moving objects. The sensor has built-in temperature compensation with high accuracy. The shell is made of stainless steel, which is waterproof and dust proof. In the same environment, our infrared temperature sensor is more stable than other products on the market.

Microphone Sensor

By using the Microphone sensor (Vocalization/Sound/Audio) we can identify the hen's Voice. By varying the abnormal sound of hen We can detect the diseases.



Fig:4.5 : Microphone sensor.

The sound sensor is one type of module used to notice the sound. Generally, this module is used to detect the intensity of sound. The applications of this module mainly include switch, security, as well as monitoring. The accuracy of this sensor can be changed for the ease of usage.

Sound Sensor Working and Its Applications

Features

These sensors are very simple to use. It gives analog o/p signal. Simply incorporates using logic modules on the input.

Specifications

The range of operating voltage is 3~5 V. The operating current is 4~5 mA. The voltage gain 26 dB ($V=6V$, $f=1kHz$). The sensitivity of the microphone (1kHz) is 52 to 48 dB. The impedance of the microphone is 2.2k Ohm. The frequency of microphone is 16 to 20 kHz. The signal to noise ratio is 54 dB.

CHAPTER 5

RESULT AND DISCUSSION:

Result

The results and discussions for early warning of poultry diseases with implications for human health would typically involve an analysis of the effectiveness of various surveillance methods, diagnostic tools, and intervention strategies in detecting and mitigating disease outbreaks.

Surveillance Methods

Present the results of surveillance efforts employed to monitor poultry populations for signs of disease. Discuss the strengths and limitations of different surveillance methods, such as passive reporting systems, active monitoring, and syndromic surveillance. Evaluate the sensitivity, specificity, and timeliness of surveillance systems in detecting disease outbreaks in poultry.

Diagnostic Tools

Describe the diagnostic tools and techniques used to identify poultry diseases, including laboratory tests, serological assays, molecular diagnostics, and imaging modalities. Present the accuracy, reliability, and practicality of diagnostic tests for early disease detection in poultry. Discuss the challenges associated with diagnosing zoonotic diseases in poultry and the implications for human health surveillance.

Intervention Strategies

Discuss the interventions implemented to control and prevent disease outbreaks in poultry populations. Evaluate the effectiveness of interventions such as vaccination programs, bio-security measures, antimicrobial stewardship, and environmental management practices. Highlight successful case studies or examples of interventions that have effectively reduced the risk of zoonotic disease transmission from poultry to humans.

One Health Approach

Emphasize the importance of a One Health approach to disease surveillance and control, recognizing the interconnectedness of human, animal, and environmental health. Discuss collaborative efforts between public health agencies, veterinary authorities, poultry producers, and other stakeholders to address poultry-related

health risks. Identify opportunities for improving communication, data sharing, and collaboration across sectors to enhance early warning systems for zoonotic disease outbreaks.

Challenges and Future Directions

Identify the challenges and gaps in current poultry disease surveillance and early warning systems. Discuss the need for improved diagnostic capabilities, enhanced data integration and analysis, and increased capacity-building efforts. Propose recommendations for future research, policy development, and public health initiatives to strengthen early warning systems for poultry diseases with implications for human health.

Conclusion

Summarize the key findings from the results and discussion section. Reiterate the importance of early disease detection in poultry for protecting human health. Highlight the significance of continued collaboration and innovation in addressing poultry-related health risks and promoting global health security. By presenting the results and engaging in discussions on these topics, researchers and policymakers can contribute to the development of effective strategies.

CHAPTER 6

SUMMARY AND CONCLUSION

In summary, early warning systems for poultry diseases play a crucial role in protecting human health by detecting and mitigating potential disease outbreaks before they escalate. Through surveillance efforts, diagnostic advancements, and intervention strategies, stakeholders aim to identify poultry diseases in their early stages, thereby minimizing the risk of zoonotic transmission to humans.

Implementing robust surveillance methods to monitor poultry populations for signs of disease, including passive reporting systems and active monitoring protocols.

Utilizing diagnostic tools and techniques to accurately identify poultry diseases, such as laboratory tests, serological assays, and molecular diagnostics.

Implementing intervention strategies, including vaccination programs, bio-security measures, and antimicrobial stewardship, to control and prevent disease outbreaks in poultry populations. Recognizing the interconnectedness of human, animal, and environmental health through a One Health approach to disease surveillance and control. Addressing challenges and gaps in current poultry disease surveillance systems, such as improving diagnostic capabilities, data integration, and collaboration across sectors.

In conclusion, early warning systems for poultry diseases are essential for safeguarding human health and promoting global health security. Continued research, collaboration, and innovation are needed to strengthen these systems, enhance early disease detection, and mitigate the risks of zoonotic disease transmission from poultry to humans.

REFERENCES

A survey of 2D shape representation: Methods, evaluations, and future research directions.

1. A. Aydin
2. A. Aydin
3. Animal Behaviour (2006).
4. Applications of Artificial Intelligence in Poultry Industry
5. Applied Animal Behaviour Science (2007)

Automatic detection of deviations in activity levels in groups of broiler chickens—a pilot study

6. Bio systems Engineering (2011)
7. Bio systems Engineering (2019)
8. Computers and Electronics in Agriculture (2014)
9. Computers and Electronics in Agriculture (2017)
10. Computers and Electronics in Agriculture (2017)

Development of an early detection system for lameness of broilers using computer vision.

Estimation of pig weight using a Microsoft Kinect prototype imaging system.

False discovery rate control is a recommended alternative to Bonferroni-type

adjustments in health studies.

Forced to crowd or choosing to cluster? Spatial distribution indicates social attraction in broiler chickens.

11. H.H. Kristensen et al

.

12. H.H. Kristensen et al.

13. J. Huang et al.

14. J. Kongsro

15. Journal of Clinical Epidemiology (2014)

16. K. Febrer et al.

17. L. Kurnianggoro et al.

18. M. Henchion et al.

19. M.E. Glickman et al.

Meat consumption: Trends and quality matters

20. Meat Science (2014)

Method for detecting avian influenza disease of chickens based on sound analysis.

The behaviour of broiler chickens in different light sources and illuminance Using 3D vision camera system to automatically assess the level of inactivity in broiler chickens.