

DEEP-AGRI: MACHINE LEARNING FOR SOIL NUTRIENTS CLASSIFICATION FOR EFFECTIVE AGRICULTURE

Submitted in partial fulfillment of the requirements for the award of
Bachelor of Engineering Degree in
Electronics and Communication Engineering

by

RAMYA A (38130182)



**DEPARTMENT OF ELECTRONICS & COMMUNICATION
ENGINEERING**

SCHOOL OF ELECTRICAL AND ELECTRONICS ENGINEERING


SATHYABAMA

**INSTITUTE OF SCIENCE AND TECHNOLOGY
(DEEMED TO BE UNIVERSITY)**

Accredited with Grade "A" by NAAC

JEPPIAAR NAGAR, RAJIV GANDHI SALAI, CHENNAI-600119

MARCH 2022



BONAFIDE CERTIFICATE

This is to certify that this Project Report is the bonafide work of **RAMYA A** (Reg. No. 38130182) who carried out the project entitled “**DEEP-AGRI: MACHINE LEARNING FOR SOIL NUTRIENTS CLASSIFICATION FOR EFFECTIVE AGRICULTURE**” under our supervision from December 2021 to March 2022.

Internal Guide

Dr. V. VIJAYAKUMAR , B. E, M.Tech,Ph.D.

Head of the Department

Dr.T. Ravi, M.E., Ph.D.,

Submitted for Viva voice Examination held on _____

Name:

Signature:

Internal Examiner

External Examiner

DECLARATION

I **RAMYA A (38130182)** hereby declare that the Project Report entitled “**DEEP-AGRI: MACHINE LEARNING FOR SOIL NUTRIENTS CLASSIFICATION FOR EFFECTIVE AGRICULTURE**” done by us under the guidance of **Dr.V.Vijayakumar** at Sathyabama Institute of Science and Technology, **Chennai- 600119** is submitted in partial fulfillment of the requirements for the award of Bachelor of Engineering degree in Electronics and Communication Engineering.

1.

2.

DATE:

PLACE:

SIGNATURE OF THE CANDIDATES

ACKNOWLEDGMENT

We are pleased to acknowledge our sincere thanks to the Board of Management of SATHYABAMA for their kind encouragement in doing this project and completing it successfully. We are grateful to them.

We convey our thanks to **Dr. N. M. NANDITHA, Dean, School of Electrical and Electronics Engineering**, and **Dr. T. RAVI, Head of the Department, Dept. of Electrical and Electronics Engineering** for providing us necessary support and details at the right time during the progressive reviews.

We would also like to express our sincere and deep sense of gratitude to our internal guide **Dr.V.VIJAYAKUMAR** , B. E, M.Tech,Ph.D. for his valuable guidance and suggestions during the progressive reviews for our project.

ABSTRACT

The major consciousness for soil management in agriculture for enhancing crop productivity is at the maintenance and development of dynamic soil parameters. The population stresses, terrestrial limitations and the decline of traditional soil management methods have directed to deterioration in the fertility of the soil in developing countries like India. Crop health is a main element within the fantastically effective gadget of cutting-edge agriculture. A substantial increase in crop production can be attained by adopting the suitable crop health management strategy.

In the existing system the research work is carried out in kerala location. The parameters such as pH value, potassium ratio, phosphorus and Nitrogen content in the soil is classified through machine learning algorithms. The proposed system focused on implementing an efficient machine learning algorithm which is nothing but the combination of more than one ML algorithm to form a hybrid network in deep learning. Deep Hybrid learning model classify the various nutrients present in the tamilnadu location, district wise and formulates a pictorial model of Location classification too. Hybrid combination of KNN and SVM Model is developed. K-nearest neighbour is the semi-supervised model and it will be more effective in parameter level classification.

The proposed model is apt for measuring the accuracy of hybrid model and readings will be tabulated.

TABLE OF CONTENTS

CHAPTER NO	TITLE	PAGE NO
1	INTRODUCTION	
	1.1 INTRODUCTION TO SOIL	6
	1.1.1 SOILS 5 INGREDIENT	6
	1.1.2 SOILS ARE BIG	6
	1.1.3 SOILS ARE YOUNG TO VERY, VERY OLD	7
	1.1.4 SOILS ARE DIVERSE	7
	1.1.5 SOILS COMMUNICATE	8
	1.1.6 SOILS ARE BIOLOGICAL BLISS	10
	1.1.7 SOILS ARE FERTILE	10
	1.1.8 SOILS ARE CLAY FACTORIES	11
	1.1.9 SOILS ARE SERVICE PROVIDERS	11
	1.2 INTRODUCTION TO SIGNAL PROCESSING TOOLBOX	11
	1.2.1 EXPLORING SIGNALS	12
	1.2.2 PREPROCESSING DATA	12
	1.2.3 FEATURE EXTRACTION AND SIGNAL MEASUREMENT SIGNAL	12
	1.3 SPECTRAL MEASUREMENT	13
	1.3.1 FILTER DESIGN AND ANALYSIS	13

	1.3.2 DESIGN FILTER	13
	1.3.3 ANALOG FILTER	13
	1.3.4 SPECTRAL ANALYSIS	13
	1.3.5 SPECTRAL ESTIMATION	13
	1.4 WINDOWS FUNCTIONS	14
	1.5 TIME FREQUENCY ANALYSIS	14
	1.5.1 TIME FREQUENCY DISTRIBUTIONS	14
	1.5.2 REASSIGNMENT AND SYNCHROSQUEEZING	14
	1.5.3 DATA ADAPTIVE TRANSFORMS	14
	1.5.4 VIBRATION ANALYSIS	14
	1.5.5 ORDER ANALYSIS	14
	1.5.6 MODEL ANALYSIS	15
	1.5.7 FATIGUE ANALYSIS	15
	1.6 ACCELERATION AND DEVELOPMENT	15
2	LITERATURE SURVEY	16
3	AIM AND SCOPE OF THE PROJECT	22
	3.1 AIM OF THE PROJECT	22

4	EXPERIMENTAL SETUP AN IMPLEMENTATION OF PROJECT	
	4.1 HARDWARE REQUIREMENTS	23
	4.2 SOFTWARE REQUIREMENTS	23
	4.2.1 MATLAB	23
	4.3 INTRODUCTION TO MATLAB 2018A	24
	4.3.1DESKTOP TOOLS AND DEVELOPMENT ENVIRONMENT	24
	4.3.2 MATHEMATICAL FUNCTION LIBRARY	24
	4.3.3 THE LANGUAGE	25
	4.3.4 GRAPHICS	25
	4.3.5 EXTERNAL INTERFACES	25
	4.4 IMAGE PROCESSING TOOLBOX	25
	4.5 COMPUTER VISION TOOLBOX	26
	4.6 IMAGE ACQUISITION	27
	4.7 RESILIENT NEURAL NETWORK	28
	4.7.1 NEURAL NETWORK	28
	4.7.2 THE BASICS OF NEURAL NETWORK	29
	4.8 GRNN NETWORK	30
	4.9 USE OF NEURAL NETWORK	32
	4.10 DESIGN METHDOLOGY	33

	4.10.1 DATA PROCESSING	34
	4.10.2 DATA CLEANING	34
	4.10.3 TRAINING DATA	35
	4.10.4 TESTING DATA	35
	4.10.5 HYBRID DEEP LEARNING NETWORK	35
	4.10.2 KNN ALGORITHM	35
5	RESULTS AND DISCUSSION	
	5.1 NEURAL NETWORKS VERSUS CONVENTIONAL COMPUTERS	36
	5.2 NETWORK LAYER	37
	5.2.1 INITIALIZE THE NETWORK	38
	5.3 TRAIN THE NETWORK	39
	5.4 TRAINING ALGORITHM	40
	5.5 INTRODUCTION CODING	41
	5.6 CODING STANDARDS	41
	5.6.1 NAMING CONVENTIONS	41
	5.6.2 VALUES CONVENTIONS	42
	5.6.3 SCRIPT WRITING AND COMMENTNG	42
	5.6.4 MESSAGE BOX FORMAT	42
6	SUMMARY AND CONCLUSION	
	6.1 CONCLUSION	48

LIST OF FIGURES

FIGURE NO	TITLE	Page no
1.1	A SOIL BODY WITH CONTRASTING AND INTERACTING MATERIALS	7
1.2	SOILS COMMUNICATE	9
4.7	NEURAL NETWORK	29
4.8	GRNN NEURAL NETWORK	31
4.10	BLOCK DIAGRAM OF DEEP-AGRI: MACHINE LEARNING FOR SOIL NUTRIENTS CLASSIFICATION FOR EFFECTIVE AGRICULTURE	34
5.2	BLOCK DIAGRAM OF NETWORK LAYERS	38
5.7	INPUT SOIL DATASET	43
5.8	PREPROCESSING	44
5.9	TRAIN THE NEURAL NETWORK	45
5.10	PATTERN RECOGNITION NEURAL NETWORK	46
5.11	SOIL ANALYSIS ACCURACY	47

CHAPTER 1

INTRODUCTION

1.1 INTRODUCTION TO SOIL:

Soils are dynamic and various natural structures that lie on the interface among earth, air, water, and lifestyles. They're critical ecosystem carrier carriers for the sustenance of humanity. The stepped forward conservation and control of soils is among the extraordinary demanding situations and opportunities we are facing inside the twenty first century.

1.1.1 SOILS 5 INGREDIENT:

Soil is a fabric composed of five substances — minerals, soil organic depend, dwelling organisms, gas, and water. Soil minerals are divided into 3 length training — clay, silt, and sand; the odds of particles in those length lessons is referred to as soil texture. The mineralogy of soils is diverse. As an example, a clay mineral known as smectite can cut back and swell a lot upon wetting and drying that it is able to knock over buildings. The maximum common mineral in soils is quartz; it makes lovely crystals but it is not very reactive. Soil organic rely is plant, animal, and microbial residues in diverse states of decomposition; it's far a important component — in fact the proportion of soil organic count in a soil is the various nice indicators of agricultural soil pleasant . Soil colorings variety from the commonplace browns, yellows, reds, grays, whites, and blacks to uncommon soil colorations together with vegetables and blues.

1.1.2 SOILS ARE BIG

You may be surprised to pay attention "dirt" defined as "huge". However, within the late 1800's soil scientists started to recognize that soils are natural bodies with size, shape, and history. Just like a water frame has water, fish, vegetation, and different components, a soil body is an included device containing soil, rocks, roots, animals, and different parts. And much like different our bodies, soil systems provide included functions which might be greater than the sum of their parts.




Fig 1.1 A soil body with contrasting and interacting materials

1.1.3 SOILS ARE YOUNG TO VERY VERY OLD

It may be difficult to mention exactly whilst a few soils had been born, however we can say that while some are young, many are very vintage. The oldest soils on this planet may be in Australia, wherein robust land office work have allowed some soils to age several million years. New soils are born with each landslide, volcanic eruption, or glacial retreat. Soils trade over time via a bunch of organic, chemical, and physical approaches. Horizons form, minerals and rocks climate, nutrients leach, and plant communities trade. Soil scientists have learned to expect the present day stage of those processes if given 5 key pieces of statistics about the soil's history — the five elements of soil formation — weather, organisms, topography, determine cloth, and time (Jenny 1941).

1.1.4 SOILS ARE DIVERSE

The sort of soil formation techniques running on one of a type decide materials under extraordinary climatic, topographic, and organic conditions over varying durations of time gives upward push to the brilliant range of soils on the earth. Soil formation creates a dizzying array of soil horizons; here are just a few examples:

- Plinthite — which hardens irreversibly upon repeated wetting and drying.
- Sulfidic — a horizon containing pyrite which, upon publicity to oxygen, can produce so much sulfuric acid that it kills plant life and may purpose fish kill.
- Petrocalcic — in which so much calcium carbonate is collected that it literally paperwork a rock-like layer in the center of a soil.

Soil scientists seize this sizeable variety thru systems of soil classification. The U.S. Machine, referred to as Soil Taxonomy, corporations soils into 12 vast orders on the most wellknown stage, and more than 19,000 soil collection on the maximum distinct degree (Ahrens & Arnold 1999). The international Union of Soil Sciences has advanced a system called the world Reference Base, which has 32 Reference Soil agencies.

1.1.5 SOILS COMMUNICATE

These grasp horizons may additionally then be in addition annotated to present extra records approximately the horizon. Horizons are first assigned to one of the following grasp horizons as distinctive through a unmarried capital letter:

O- Horizon containing a excessive percentage of soil organic be counted.

- A - Horizon darkened by using the buildup of natural rely.
- E - Horizon shaped through the elimination (eluviations) of clays, natural remember, iron, or aluminum. Commonly lightened in shade due to those removals.
- B - extensive elegance used for subsurface horizons which have been transformed drastically by using a soil formation process inclusive of colour and shape improvement; the deposition (illuviation) of substances inclusive of clays, natural depend, iron, aluminum, carbonates, or gypsum; carbonate or gypsum loss; brittleness and excessive density; or excessive weathering main to the buildup of weathering-resistant minerals.

- C - A horizon minimally affected or unaffected with the aid of the soil formation techniques.
- R - Bedrock.

Those grasp horizons may additionally then be in addition annotated to provide additional records approximately the horizon. Decrease case letters can be located as suffixes following the master horizon letter to present additional records about soil characteristics or soil formation approaches. As an instance, the lower case "t" on the B horizon in shows that the horizon is characterized with the aid of illuvial clay accumulation. Multiple letters can be used — depicts a Bkm horizon that means that it's miles cemented (m) by way of illuvial carbonates (ok). Numbers located before the grasp horizon name (e.g., 2Bt) indicate a difference in discern fabric; numbers positioned on the cease of a horizon name are used to subdivide horizons that have the identical designation however are exceptional in some manner (e.g., a crimson Bt1 over a yellow Bt2). Example: soil with designations that communicate the soil formation processes occurring in each horizon.




Fig 1.2 soils communicate

1.1.6 SOILS ARE BIOLOGICAL BLISS

If you want existence, you will love soils. There are a host of small, medium, and massive organisms that live in soils, including mammals, birds, insects, and protozoa. However the finest biodiversity lies inside the soil microbes — the bacteria, fungi, and archaea. A teaspoon of wealthy soil can incorporate a billion micro organism. We truly realize very little about the diversity of soil microbes, partly due to the fact they may be so numerous, but also because we've got now not been capable of tradition the full-size majority of these organisms inside the lab. Soil microbiologists are making use of superior molecular techniques to apprehend the variety and feature of soil microbes. It is an exciting field of exploration with new biological taxa regularly being found.

1.1.7 SOILS ARE FERTILE

Soils are the number one provider of vitamins and water for a great deal of the flowers in the world. There are 18 factors considered vital for plant boom, maximum of which might be made available to flowers thru root uptake from soils (Brady & Weil 2007). Soils keep nutrients by way of several mechanisms. Maximum vitamins are dissolved in soil water as either undoubtedly or negatively charged ions; soil debris also are charged and thereby are capable of electrically maintain these ions. Soils also keep nutrients through preserving the soil water itself. Arguably the finest of all of the surroundings offerings supplied by soils is the retention of water — with out soils our land could be little however rocky deserts. Flowers use tons greater water than one may assume because they're constantly releasing water into the atmosphere as a result of transpiration, which is a component of the system of photosynthesis. Clay and silt particles are the primary mineral components in soils that preserve water — those small particles gradual the drainage of water and, like a sponge, physically hold water via capillary forces. Clay provides such strong force that plant life can't pull all the water away from it, which makes silt particles the final element for plant-to be had water storage — they maintain large portions of water but additionally launch it to plant roots.

1.1.8 SOILS ARE CLAY FACTORIES

Most of the maximum crucial capabilities accomplished via soils is to provide the perfect conditions for clay synthesis. Clays are vital because they are often lively, that is a preferred term soil scientists use to explain how chemically reactive a particle is with ions, water, and other particles. These reactions are critical for the provision of many environment offerings. Clays are often the maximum energetic mineral particles due to the fact they have got particular chemical characteristics and also due to the fact they have got so much surface vicinity — clays will have 10,000 instances the floor location of sand of the identical weight (Brady & Weil 2007). All this surface place makes clays a warm spot for chemical reactions.

1.1.9 SOILS ARE SERVICE PROVIDER

Soils are the among the extremely good atmosphere service companies on earth (Haygarth & Ritz 2009). They keep and provide water for plants. They prevent floods with the aid of transferring water slowly to streams and groundwater. They clear out and remediate pollution. They cycle and recycle nutrients and wastes — transforming them into biologically to be had paperwork, storing them away for later use, and stopping their leaching to floor and floor waters. Soils provide habitat for a full- size range of life. They take up and launch crucial gases, including oxygen and greenhouse gases, a provider called gasoline regulation. Lots of those atmosphere offerings are being lost via the degradation and loss of soils. The conservation, recovery, and optimization of environment services furnished by using soils is most of the outstanding demanding situations for humanity inside the twenty first century.

1.2 INTRODUCTION TO SIGNAL PROCESSING TOOLBOX

Matlab has provided many fundamental signal and device processing functions which includes convolution, Fourier transform, Laplace transform, z-remodel, and many others, which simplify the calculation method significantly. Matlab offer numerous strategies to analyze the indicators and systems, including both non-stop and discrete situations.

1.2.1 EXPLORING SIGNALS

Use the sign Analyzer app to analyze and visualize indicators inside the time, frequency, and time-frequency domains. Extract areas of interest from alerts for in addition evaluation.

The signal Analyzer app also permits you to degree and examine indicators of various durations on the same time and within the same view.

1.2.2 PREPROCESSING DATA

Denoise, clean, and detrend indicators to prepare them for similarly evaluation. Do away with outliers and spurious content from records.

Decorate indicators, visualize them, and find out styles. Trade the pattern fee of a sign or make the sample charge constant for irregularly sampled indicators or indicators with lacking statistics.

1.2.3 FEATURE EXTRACTION AND SIGNAL MEASUREMENTS

- Measure not unusual different features and extract styles in alerts.

Descriptive records

- Compute commonplace descriptive facts like maxima, minima, wellknown deviations, and RMS ranges. Find changepoints in indicators and align alerts the use of dynamic time warping.
- Find sign peaks and determine their top, width, and distance to buddies. Degree time-area capabilities including top-to-height amplitudes and sign envelopes.
- Pulse and Transition Metrics

Measure upward push time, fall time, slew fee, overshoot, undershoot, settling time, pulse width, pulse length, and duty cycle.

1.3 SPECTRAL MEASUREMENTS

Compute the bandwidth and imply or median frequency for alerts or energy spectrum. Degree signal-to-noise ratio (SNR), general harmonic distortion (THD), and signal-to-noise and distortion ratio (SINAD). Degree harmonic distortion. Estimate immediate frequency, spectral entropy, and spectral kurtosis.

1.3.1 FILTER DESIGN AND ANALYSIS

Layout, examine, and put in force a variety of virtual and analog filters.

1.3.2 DIGITAL FILTER

Layout, examine, and put in force a spread of virtual FIR and IIR filters, inclusive of lowpass, highpass, and bandstop, using the filter dressmaker app. Visualize importance, section, institution put off, impulse, and step responses.

Study clear out poles and zeros. Examine clear out performance with the aid of checking out stability and phase linearity. Observe filters to facts and eliminate delays and segment distortion using zero-segment filtering.

1.3.3 ANALOG FILTER

Layout and examine analog filters, consisting of Butterworth, Chebyshev, Bessel, and elliptic designs. Perform analog-to-virtual clear out conversion the usage of discretization strategies including impulse invariance and the bilinear transformation.

1.3.4 SPECTRAL FILTER

Characterize the frequency content of a signal.

1.3.5 SPECTRAL ESTIMATION

Estimate spectral density using nonparametric strategies along with the periodogram, Welch's overlapped phase averaging method, and the multitaper method. Put in force parametric and subspace techniques such as Burg's, covariance, and track to estimate spectra. Compute electricity spectra of non

uniformly sampled signals or indicators with lacking samples using the Lomb-Scargle method. Degree sign similarities inside the frequency area by means of estimating spectral coherence.

1.4 WINDOWS FUNCTIONS

Implement and visualize not unusual window features. Use the Window designer app to layout and examine home windows. Compare main lobe widths and side lobe degrees of home windows as a characteristic in their length and other parameters.

1.5 TIME FREQUENCY ANALYSIS

Visualize and examine time-frequency content of nonstationary alerts.

1.5.1 TIME FREQUENCY DISTRIBUTIONS

Use the quick-time Fourier remodel, spectrograms, or Wigner-Ville distributions to research indicators with time-various spectral content. Use the move spectrogram to compare signals inside the time-frequency area.

1.5.2 REASSIGNMENT AND SYNCHROSQUEEZING

Use the reassignment approach to sharpen the localization of time-frequency estimates. Discover time-frequency ridges using synchro squeezing

1.5.3 DATA ADAPTIVE TRANSFORMS

Carry out facts-adaptive time-frequency evaluation the usage of empirical mode decomposition, variational mode decomposition and Hilbert-Huang remodel

1.5.4 VIBRATION ANALYSIS

Characterize vibrations in mechanical systems.

1.5.5 ORDER ANALYSIS

Use order analysis to investigate and visualize spectral content taking place in rotating machinery.

Track and extract orders and their time-area waveforms. Tune and extract RPM profiles from vibration alerts. Remove noise coherently with time-synchronous averaging.

1.5.6 MODEL ANALYSIS

Perform experimental modal evaluation by way of estimating frequency-response functions, natural frequencies, damping ratios, and mode shapes.

1.5.7 FATIGUE ANALYSIS

Generate excessive-cycle rainflow counts for fatigue evaluation.

1.6 ACCELERATION AND DEPLOYMENT

Use GPUs to accelerate the code. Generate portable C/C++ supply code, standalone executables, or standalone applications from your MATLAB® code.

Accelerating the Code

Speed up the code with the aid of using GPU and multicore processors for supported capabilities.

CHAPTER 2

LITERATURE SURVEY

X. C. Cui and J. L. Hu (2013) [1] ,The nitrogen (N) nutrients, crop yield, and responses of wheat to arbuscular mycorrhizal fungi (AMF) had been tested in an experimental area under unfastened-air ozone concentration [O₃] enrichment (FACE) situations. The test included three Treatments: ambient [O₃] (Ambient), increased [O₃] (FACE, centered at ambient [O₃] $\times$ 1.5), and accelerated [O₃] inoculated with an AMF consortium which include numerous *Glomus* species (FACE+AMF). AMF inoculation responsiveness of wheat became predicted through evaluating flowers grown in unsterilized soil inoculated with the exogenous AMF and in untreated soil containing indigenous AMF. As compared with the Ambient, fairly higher N contents however lower shoot biomasses of wheat plants had been observed within the FACE remedy without AMF inoculation from the tillering degree in February and heading degree in April, respectively, which appreciably ($P < 0.05$) decreased grain yield by means of 28% at harvest in June. Beneath the FACE situation, as compared with the non-inoculated remedy, AMF inoculation considerably ($P < 0.05$) extended root colonization charges each at the tillering degree and heading degree, and also considerably ($P < 0.05$) multiplied shoot biomass at the heading stage and, hence, appreciably ($P < 0.05$) increased grain yield by using 40% at harvest. But, AMF inoculation considerably ($P < 0.05$) reduced total N content material in wheat shoots at the tillering level, suggesting that AMF consortia can also beautify plant tolerance to expanded [O₃] through elevating root colonization price in preference to plant overall N content at early growing levels.

Fuji Jian Digvir S. Jayas Noel D. G. White (2013) [2], Ozone (O₃) is a strong oxidant with an extended history of safe use in lots of fields. Researches have been lately focused at the utility of O₃ as a fumigant to manipulate saved-grain bugs and microorganisms and to lessen mycotoxins. This evaluate located the following records: (1) O₃ substantially suppressed insect populations at B50 ppm with four days treatment; (2) to eliminate insect infestation,[a 135 ppm with greater than 8 days remedy could be required; (3) O₃ at 50 ppm with 3 days treatment decreased 63 % of stored fungi; (4) O₃ at 5–30 ppm could reduce mycotoxin

contamination; however, high awareness and long remedy time had been required to remove mycotoxins; (5) utility of O₃ at doses that were sufficient for the effective disinfection of grain might have an effect on characteristics of grain; and (6) O₃ at 47–106 ppm should noticeably harm gadget in 2 months with the aid of corrosion. Based on those information, we advocated that B50 ppm O₃ have to be used within the saved-grain enterprise and its capability method of utility became also analyzed.

M.Nur, E.Kusdiyantini, W.Wuryanti(2015) [3], This research has been carried out by using the use of ozone to address the rapidly declining first-class of rice in garage. Inside the first 12 months, research has focused on the rice storage with ozone generation for small ability (e.g., household) and the medium ability (e.g., dormitories, hospitals). Ozone become produced with the aid of an ozone generator with Dielectric Barrier Discharge Plasma (DBDP). Ozone generation rice storage system (OTRISS) is using ozone characteristic which is a sturdy oxidizer. Ozone have a short staying power of existence and then decompose, as a end result produce oxygen and radicals of oxygen. These characteristics should kill micro-organisms and pests, reduce air humidity and improve oxygen. All additives used in SPBTO assembled the usage of raw materials available in the massive towns in Indonesia. Provider of excessive voltage (excessive Voltage strength supply, 40-70 kV, 23 KH, AC) is one of components which have been assembled and examined. Ozone generator is assembled with 7 reactors of Dielectric Barrier Discharge Plasma (DBDP). Rice field which have been prepared for OTRISS have adjusted so can be incorporated with generator, energy deliver and blower to blow air. OTRISS with a potential of 75 kg and 100 kg have been made and tested. The potential of ozone to cast off bacteria and fungi have been examined and resulted in a decrease of microorganisms at 3 log CFU/g. Checking out in food chemistry showed that ozone treatment of rice had now not changed the chemical content that also meet the usual of chemical content and dietary applicable to ISO wellknown milled rice. The results of this examine are very probably to be used as an opportunity to rice garage structures in warehouse. Check and scale-up is being achieved in a mini warehouse whose circumstance is mimicked to rice in national Rice storage of Indonesia to ensure excellent. Next diversifications could be set up within the rice storage system within the Bulog.

Xuelian Bao Qi Li Jianfeng Hua (2014) [4], Ultraviolet-B (UV-B) radiation and increased tropospheric ozone may additionally motive reductions inside the productivity and high-quality of crucial agricultural vegetation. But, research concerning their interactive impact remains scarce, in particular on the belowground techniques. The use of the open top chambers experimental setup, we monitored the reaction of soil nematodes to the increased O₃ and UV-B radiation personally in addition to in aggregate. Our consequences indicated that increased O₃ and UV-B radiation have effect not best on the belowground biomass of flora, but additionally on the network shape and purposeful range of soil nematodes. The canonical correspondence analysis suggested that soil pH, shoot biomass and microbial biomass C and N had been relevant parameters that influencing soil nematode distribution. The interactive outcomes of expanded O₃ and UV-B radiation was best located on the abundance of bacterivores. UV-B radiation appreciably expanded the abundance of overall nematodes and bacterivores in evaluation with the manipulate at pod-filling degree of soybean. Following elevated O₃, nematode range index reduced and dominance index elevated relative to the manipulate at pod-filling level of soybean. Nematode functional diversity showed response to the results of increased O₃ and UV-B radiation at pod-bearing degree. Higher enrichment index and lower structure index within the treatment with both increased O₃ and UV-B radiation indicated a confused soil circumstance and degraded soil meals internet. However, The ratios of nematode trophic organizations cautioned that the negative outcomes of accelerated O₃ on soil meals internet may be weakened through the UV-B radiations.

Whangchai, K Uthaibutra (2016) [5], Ethion is an organophosphate pesticide which is widely used in tangerine cultivation. This studies become performed as an research at the efficiency of ozone microbubbles and ultrasonic irradiation in mixture (OMB/US) on ethion residue discount in tangerine cv. Sai Nam Pung. Ethion preferred (10 ppm) dealt with with OMB/US become as compared with individual ozone microbubbles (OMB) and ultrasonic frequency 1000 kHz (US) at various time intervals (15, 30, 45, and 60 min). It was discovered that the ethion awareness of dealt with samples had considerably decreased in direct percentage to the response time, compared with the manage. The evaluation was finished to

find out which OMB/US remedy confirmed decrease case heading/subheading the most effectiveness on ethion degradation over OMB or US alone. The rate steady (okay) of OMB/US changed into revealed as $0.79 \times 10^{-2} \text{ min}^{-1}$, that is the maximum value obtained of all of the remedies. The detoxification stages of OMB/US, OMB, and US had been determined the use of the brine shrimp lethality take a look at technique. OMB/US confirmed the maximum degree of LC50, at 281.43 mgL⁻¹, whilst compared the manage which have LC50 at 5.41 mgL⁻¹, the ethion residue on tangerine fruit turned into treated via OMB/US, the same as the previous technique. The outcomes confirmed that the percentage of ethion degradation had accelerated, which turned into similarly to the upward thrust inside the response time. OMB/US confirmed the highest fee of discount as occurring inside 15 min. It become located that ethion had reduced with the aid of 73% after OMB/US treatment for 60 min reaction time. Thereafter, tangerine fruits had been saved at 5°C for 35 days to determine excellent adjustments. The peel color exchange, percent of weight reduction, TSS, TA, and ascorbic acid content after all of the treatments had been found to be not stricken by OMB/US.

Ebihara, F. Mitsugi(2014) [6], We've developed an ozone-mist spray gadget for chemical unfastened agricultural sterilization. This environmentally friendly gadget is the movable and backpack kind wherein the excessive density ozone is generated with the aid of surface dielectric barrier discharge in oxygen. The ozone fuel is injected into the water-mist that's produced by means of the water spray nozzle. The ozone-mist is sprayed over dangerous bugs, viruses and micro organism living on/inside the flora. The ozone solubility of the ozone-mist water is over 5ppm in the case of wellknown treatment situations (oxygen flow charge 1 L/min, mist water go with the flow charge 0.3 L/min). Many sorts of dangerous insects have been sterilized to evaluate the sterilization characteristics and operation management through changing conditions together with ozone concentration, oxygen float charge, mist water flow charge, distance among nozzle and organic gadgets the usage of various structures of the spray nozzle machine. Ordinary experiment indicates that the survival charge in the case of aphids (small bugs of a few mm size) turns into sensible value of approximately 10% for very quick spray time of 10 sec.

Hao Zhang, Danyan Ma (2016) [7], There is an increasing focus round the world on soil infection through natural compounds because of its harmful impact on human health. Considerable efforts had been committed to the remediation of soil infected with the aid of organics, leading to the development of various remediation techniques, together with physical remediation, bioremediation, and chemical remediation. Currently, an appealing superior oxidation process (AOP), non-thermal plasma technology, has drawn increasing interest for the remediation of natural contaminated soil due to the low strength consumption, rapid begin-up and shutdown of the system, in addition to a low requirement for the pretreatment procedure of soil, etc. In this paintings, the present day state of affairs of soil contamination by using natural pollutants and the existing soil remediation methods have been delivered. The studies progresses of numerous non-thermal plasmas for organic infected soil remediation had been offered, consisting of dielectric barrier discharge (DBD), pulsed corona plasma, and non-thermal gliding arc fluidized bed. The results of various parameters (e.g., carried out voltage, soil homes, kind and feed drift fee of the service gasoline, in addition to the reactor configuration) at the remediation performance of nonthermal plasmas had been discussed. In addition, the viable degradation mechanisms of natural pollutants . Via non-thermal plasmas were analyzed.

Chakraborty, Subhra; Salekdeh, Ghasem Hosseini (2015) [8], Within the hastily developing economies of Asia and Oceania, food safety has come to be a primary issue. With the rising populace, growing greater meals at low cost prices is turning into even extra crucial. Similarly, the expected weather exchange will lead to drastic adjustments in global floor temperature and adjustments in rainfall patterns that during flip could pose a extreme danger to plant vegetation worldwide. As a end result, understanding how plant life will live to tell the tale in a converting weather will be increasingly crucial. Such challenges require included tactics to growth agricultural production and deal with environmental threats. Proteomics can play a role in resolve the underlying mechanisms for meals manufacturing to deal with the developing call for for meals. On this evaluation, the current fame of food crop proteomics is mentioned, particularly in regards to the Asia and Oceania areas. Furthermore, the destiny attitude in terms of proteomic techniques for the crucial meals crops is highlighted.

Takamasa SAKAI, Joanna PAWLAT (2012) [9], We have studied gaseous ozone sterilization for soil and biomedical prevention for infectious illnesses and for the protective of citizens' health. Experimental consequences received the usage of ozone era apparatuses of numerous production and ozone sterilization structures, following with AFM dimension were proven. Alternate of soil properties within the method of ozone remedy, plant increase, biological phenomena and DNA visualization were investigated. Utility of this method to environment safety become proposed.

Nahed Msayleba, Saiid Ibrahim (2011) [10], This have a look at tests Ozone as a Nematicides' alternative. Nematode-infected soil samples had been dealt with with ascending doses of O₃ with the aid of submerging the outlet of an "MB1000 Ozone Generator" within the 40 ml samples; then to test the O₃ nematicidal effect by fumigation, Ozone gas was released right into a sealed bag containing 80g of each of the 6 nematode-infected soil samples with ascending doses and a repetition of every. With water-ozonation, 900 mg O₃ have been had to kill 100% of nematode, and the O₃-Nematodes LD₅₀ turned into diagnosed through 420mg. With the second one test, O₃ soil fumigation for 50 mins at a dose of 1,125 mg in an air volume of five liters, were needed to control 95% of dwelling nematodes.

CHAPTER 3

AIM AND SCOPE OF THE PROJECT

The main reason for a heavy loss in soil quality is due to incorrect soil and crop management strategies. Also excess use of chemical fertilizers has created imbalances in the availability of soil nutrients. In the current agricultural scenario, an early prediction about the soil status is very helpful for farmers to create a better environment for next cultivation. So machine learning techniques help in the domain of agriculture and could support data analysis for making prediction.

3.1 AIM OF THE PROJECT

Hence the classification of the ph and relevant soil nutrients indices help to save the time of analysis soil health and environmental quality. Hybrid ml algorithm is implemented to improve the accuracy and for multi-feature soil resourced particular location. The proposed system focused on implementing an efficient machine learning algorithm which is nothing but the combination of more than one ml algorithm to form a hybrid network in deep learning. Deep hybrid learning model classify the various nutrients present in the tamilnadu location, district wise and formulates a pictorial model of location classification too.

From the literature studies the objective is confined to,

- Save the time of analysis soil health
- Low computation time
- Performs well with large dataset
- Easy to implement

CHAPTER 4

EXPERIMENTAL SETUP AND IMPLEMENTATION OF PROJECT

4.1 HARDWARE REQUIREMENTS

- Hard Disk : 500GB and Above
- RAM : 4 GB and Above
- Processor : i3 and Above

4.2 SOFTWARE REQUIREMENTS

- MATLAB R2018a
- Windows Operating System XP and Above

4.2.1 MATLAB

Matlab (matrix laboratory) is a multi-paradigm numerical computing surroundings and fourth-technology programming language. A proprietary programming language developed by way of Math Works, MATLAB permits matrix manipulations, plotting of functions and information, implementation of algorithms, advent of consumer interfaces, and interfacing with applications written in different languages, which includes C, C++, Java, Fortran and Python.

Despite the fact that MATLAB is intended ordinarily for numerical computing, an optionally available toolbox makes use of the Mu PAD symbolic engine, allowing access to symbolic computing abilities. An additional bundle, Simulink, provides graphical multi-domain simulation and model-based design for dynamic and embedded structures.

The basic statistics shape in MATLAB is the array, an ordered set of real or complex elements. This object is evidently ideal to the representation of pictures, actual-valued ordered sets of shade or intensity data.

MATLAB shops most images as -dimensional arrays (i.e., matrices), in which each detail of the matrix corresponds to a unmarried pixel within the displayed image. (Pixel is derived from image element and commonly denotes a single dot on a

laptop display.) as an example, an picture composed of 200 rows and 300 columns of various colored dots could be saved in MATLAB as a two hundred-via-three hundred matrix. A few photos, which include RGB, require a three-dimensional array, where the first plane inside the 0.33 size represents the crimson pixel intensities, the second one plane represents the green pixel intensities, and the third aircraft represents the blue pixel intensities.

This conference makes running with snap shots in MATLAB similar to working with some other type of matrix facts, and makes the overall energy of MATLAB available for image processing packages. As an example, you could pick out a single pixel from an image matrix the usage of normal matrix subscripting.

4.3 INTRODUCTION TO MATLAB R2018a

The MATLAB device consists of those primary components.

4.3.1 Desktop Tools and Development Environment

This is the set of equipment and facilities that help you operate and become greater effective with MATLAB features and files. A lot of these gear are graphical consumer interfaces. It includes the MATLAB computing device and Command Window, a command records, an editor and debugger, a code analyzer and different reports, and browsers for viewing help, the workspace, documents, and the hunt path.

4.3.2 MATHEMATICAL FUNCTION LIBRARY

That is a extensive series of computational algorithms starting from standard functions, like sum, sine, cosine, and complex mathematics, to greater state-of-the-art functions like matrix inverse, matrix Eigen values, Bessel capabilities, and fast Fourier transforms.

4.3.3 THIS LANGUAGE

This is a high-degree matrix/array language with manipulate go with the flow statements, functions, facts structures, enter/output, and item-oriented programming capabilities. It lets in both "programming in the small" to rapidly create quick and grimy throw-away programs, and "programming within the massive" to create huge and complicated application packages.

4.3.4 GRAPHICS

MATLAB has massive facilities for showing vectors and matrices as graphs, in addition to annotating and printing those graphs. It includes high-level functions for 2-dimensional and 3-dimensional facts visualization, image processing, animation, and presentation pix. It's also low-stage functions that can help you completely customize the advent of photos as well as to construct whole graphical user interfaces to your MATLAB packages.

4.3.5 EXTERNAL INTERFACE

That is a library that permits you to jot down C and FORTRAN programs that interact with MATLAB. It includes centers for calling routines from MATLAB, for calling MATLAB as a computational engine, and for reading and writing MAT-files.

4.4 IMAGE PROCESSING TOOLBOX

The image Processing Toolbox is a collection of features that enlarge the capability of the MATLAB numeric computing surroundings. The toolbox helps a extensive range of photo processing operations, such as:

- Spatial photograph transformations
- Morphological operations
- community and block operations
- Linear filtering and filter layout
- Transforms
- picture analysis and enhancement

- photograph registration
- De blurring
- region of interest operations

Many of the toolbox features are MATLAB M-documents, a sequence of MATLAB statements that put in force specialized photograph processing algorithms. You may increase the skills of the photo Processing Toolbox by using writing your own M-files, or by means of the use of the toolbox in aggregate with other toolboxes, inclusive of the signal Processing Toolbox and the Wavelet Toolbox.

4.5 COMPUTER VISION TOOLBOX

The sphere is of significance for such numerous programs as autonomous cars, navigating with the help of images, captured with the aid of a mounted digicam, and excessive precision measurements using photographs, taken with the aid of calibrated cameras. In this paper, we will gift some of numerical routines, applied in MATLAB, that are useful in a variety of computer imaginative and prescient applications. The collection of routines can be called the laptop vision Toolbox. One of the fundamental problems in laptop imaginative and prescient is to calculate the three-D-shape of the scene and the motion of the camera from measurements inside the snap shots taken from exceptional view-points inside the scene. This hassle is called structure and movement, relating to the reality that both the structure of the scene and the motion of the digital camera are calculated from photograph measurements best. Some of special sub issues, springing up from one-of-a-kind know-how of the intrinsic residences of the camera seem. Other vital troubles are to calculate the shape of the 3D-scene given the movement of the digital camera and to calculate the motion of the digicam given the structure of the scene. These problems are incredibly less difficult than the overall shape and movement problem, but are though important for navigation and obstacle avoidance. A related trouble is to calibrate a camera, i.e. Calculate the focal distance, the primary point and many others., from photos of a known scene. This calibration may be used to make a extra unique reconstruction. The problem of estimating shape and motion from image sequences are intently associated with the field of closed variety photogrammetric. In this field photos are used to make

specific measurements of three-dimensional objects from some of photographs, often taken with the aid of calibrated cameras. The primary difference is that during computer vision, the focal point isn't always on accuracy, however as an alternative on reliability and velocity of calculation. In these kind of troubles different type of so called capabilities may be used. The only function is factors which are easily detected in the sequence, inclusive of corners or pain- ted marks. Other types of capabilities are strains and curves, which can be less difficult to locate and track but extra hard to use in structure and motion algorithms. If you want to use our laptop vision exercises these functions ought to be detected in advance and the correspondence between different pictures ought to be established. The main emphasis of our computer vision toolbox is to apply these detected functions to resolve for shape and motion.

4.6 IMAGE ACQUISITION TOOLBOX

The image Acquisition Toolbox's is a group of functions that expand the functionality of the MATLAB numeric computing environment. The toolbox helps a wide variety of picture acquisition operations along with

- acquiring images through many types of photo acquisition gadgets, from professional grade frame grabbers to USB-primarily based Webcams
- Viewing a preview of the stay video movement
- Triggering acquisitions (includes outside hardware triggers)
- Configuring call lower back capabilities that execute while certain events arise
- Bringing the picture records into the MATLAB workspace

Among the toolbox features are MATLAB M-files. You may make bigger the abilities of the photograph Acquisition Toolbox with the aid of writing your personal M-documents, or by means of the use of the toolbox in combination with other toolboxes, consisting of the picture Processing Toolbox and the data Acquisition Toolbox. The toolbox also includes a Simulink interface referred to as the photo

Acquisition Block set. This block set extends Simulink with a block that helps you to convey live video statistics into a version.

4.7 RESILIENT NEURAL NETWORK:

Resilient returned propagation (Rprop), an set of rules that may be used to teach a neural network, is similar to the extra not unusual (ordinary) lower back-propagation. However it has foremost benefits over lower back propagation: First, training with Rprop is often quicker than education with lower back propagation

4.7.1 NEURAL NETWORK

The most common form of artificial neural community includes three businesses, or layers, of devices: a layer of "input" gadgets is attached to a layer of "hidden" devices, that is connected to a layer of "output" devices.

The pastime of the enter units represents the raw data this is fed into the community.

The activity of every hidden unit is determined by using the activities of the enter devices and the weights at the connections among the enter and the hidden devices.

The behavior of the output devices depends on the interest of the hidden gadgets and the weights between the hidden and output gadgets.

This simple kind of community is interesting because the hidden devices are loose to assemble their very own representations of the enter. The weights among the input and hidden devices determine while every hidden unit is energetic, and so through modifying those weights, a hidden unit can select what it represents.

We additionally distinguish single-layer and multi-layer architectures. The unmarried-layer company, in which all devices are connected to each other, constitutes the most standard case and is of greater capacity computational energy than hierarchically established multi-layer corporations. In multi-layer networks, gadgets are often numbered by using layer, in preference to following a global numbering. A Neural network (NN) is an facts processing paradigm that is stimulated by using the manner organic apprehensive systems, inclusive of the

mind, procedure statistics. The key detail of this paradigm is the novel structure of the facts processing machine. It's miles composed of a big quantity of relatively interconnected processing elements (neurons) operating in unison to solve specific issues. NNs, like people, analyze by means of example. An NN is configured for a specific software, such as sample popularity or data type, through a getting to know technique. Learning in organic structures includes changes to the synaptic connections that exist among the neurons.

4.7.2 THE BASICS OF NEURAL NETWORKS

Neural networks are usually organized in layers. Layers are made from a number of interconnected 'nodes' which include an 'activation function'. Styles are supplied to the network thru the 'enter layer', which communicates to 1 or greater 'hidden layers' where the actual processing is achieved through a device of weighted 'connections'. The hidden layers then hyperlink to an 'output layer' in which the solution is output as proven within the photo underneath.




Fig 4.7 neural network

All NN network has 4 layers:

- **Input layer** — there's one neuron inside the input layer for every predictor variable. In the case of specific variables, N-1 neurons are used wherein N is the quantity of classes. The enter neurons (or processing earlier than the input layer) standardizes the variety of the values with the aid of subtracting

the median and dividing by using the interquartile variety. The input neurons then feed the values to each of the neurons within the hidden layer.

- **Hidden layer** — this residue has one neuron for every case within the schooling statistics set. The neuron shops the values of the predictor variables for the case in conjunction with the target value. While offered with the x vector of input values from the input layer, a hidden neuron computes the Euclidean distance of the test case from the neuron's middle point and then applies the RBF kernel feature the usage of the sigma value(s). The ensuing cost is passed to the neurons within the sample layer.
- **Sample layer / Summation layer** — the subsequent layer within the community is one-of-a-kind for NN networks and for GRNN networks. For NN networks there may be one sample neuron for each category of the goal variable. The real goal category of every training case is stored with every hidden neuron; the weighted price popping out of a hidden neuron is fed handiest to the sample neuron that corresponds to the hidden neuron's class. The pattern neurons add the values for the magnificence they constitute (consequently, it's miles a weighted vote for that class).
- **Decision layer** — The decision layer is exceptional for NN and GRNN networks. For NN networks, the selection layer compares the weighted votes for every goal class accrued inside the pattern layer and uses the most important vote to expect the target category.

4.8 GRNN NETWORK

For GRNN networks, the decision layer divides the cost collected within the numerator summation unit via the value inside the denominator summation unit and makes use of the result because the expected goal fee.




Fig 4.8 GRNN NEURAL NETWORK

- **Input Layer**--The enter vector, denoted as p , is presented as the black vertical bar. Its measurement is $R \times 1$. On this paper, $R = 3$.
- **Radial foundation Layer**-- In Radial foundation Layer, the vector distances between enter vector p and the load vector made of every row of weight matrix W are calculated. Right here, the vector distance is described as the dot product between vectors [8]. Expect the size of W is $Q \times R$. The dot product between p and the i -th row of W produces the i -th detail of the space, whose size is $Q \times 1$. The minus image, “-”, suggests that it is the space between vectors. Then, the bias vector b is mixed with the aid of an detail-by-element multiplication, .The end resulttransfer characteristic in NN has built into a distance criterion with appreciate to a center. In this paper, it is defined as $\text{radbas}(n) = 2 n e^{- (1)}$ each element of n is substituted into Eq. 1 and produces corresponding detail of a , the output vector of Radial foundation Layer. The i -th element of a may be $..B_i$ (2) where W_i is the vector made of the i -th row of W and b_i is the i -th detail of bias vector b .
- **A few traits of Radial basis Layer**--The i -th detail of a equals to one if the enter p is same to the i th row of enter weight matrix W . A radial foundation neuron with a weight vector close to the input vector p produces a cost close to 1 and then its output weights inside the aggressive layer will pass their values to the competitive characteristic. It is also feasible that numerous factors of a are close to 1 since the input pattern is near several training styles.

- **Competitive layer**--there is no bias in competitive Layer. In aggressive Layer, the vector a is first of all elevated with layer weight matrix M , producing an output vector d . The Competitive function, denoted as C in Fig. 2, produces a 1 corresponding to the most important element of d , and 0's some other place. The output vector of aggressive function is denoted as c . The index of 1 in c is the quantity of tumor that the device can classify. The dimension of output vector, ok , is five in this paper.

4.9 USE OF NEURAL NETWORKS:

Neural networks, with their high-quality capacity to derive that means from complex or vague facts, can be used to extract patterns and stumble on tendencies that are too complicated to be observed through either people or different laptop strategies. A educated neural community can be concept of as an "expert" in the category of records it has been given to research. This expert can then be used to offer projections given new situations of interest and answer "what if" questions.

Different blessings include:

1. Adaptive studying: An potential to discover ways to do duties based totally on the facts given for training or preliminary revel in.
2. Self-organisation: An NN can create its own organization or representation of the statistics it gets during learning time.
3. Real Time Operation: NN computations can be completed in parallel, and unique hardware devices are being designed and synthetic which take gain of this functionality.
4. Fault Tolerance via Redundant statistics Coding: Partial destruction of a community results in the corresponding degradation of overall performance. However, some community competencies may be retained inspite of main network damage.

4.10 DESIGN METHODOLOGY

In the existing system the research work is carried out in kerala location. The parameters such as ph value, potassium ratio, phosphorus and Nitrogen content in the soil is classified through machine learning algorithms. The proposed system focused on implementing an efficient machine learning algorithm which is nothing but the combination of more than one ML algorithm to form a hybrid network in deep learning. Deep Hybrid learning model classify the various nutrients present in the tamilnadu location, district wise and formulates a pictorial model of Location classification too. Hybrid combination of KNN and SVM Model is developed. K-nearest neighbor is the semi-supervised model and it will be more effective in parameter level classification. The proposed model is apt for measuring the accuracy of hybrid model and readings will be tabulated. Hybrid ML algorithm is implemented to improve the accuracy and for multi-feature soil resourced tamil-nadu location




Fig 4.10 block diagram of deep-agri: machine learning for soil nutrients classification for effective agriculture

4.10.1 DATA PREPROCESSING

This module is used to read the soil dataset, prepare it to make a readable format, labelize the values, plot the required histograms for analysis etc.

4.10.2 DATA CLEANING

This module is used to remove unwanted numbers and empty data.

4.10.3 TRAINING DATA

It is the initial data used to train machine learning model. This data set are fed to machine learning algorithm to teach them how to make prediction or perform a desired task.

4.10.4 TESTING DATA

This module is used to evaluate models accuracy. It make prediction when processing unseen data.

4.10.5 HYBRID DEEP LEARNING NETWORK

. Hybrid deep learning network is a combination of knn and hybrid classification and deep learning network. This module is used to implement a hybrid classification model to determine the soil nutrients present in particular location and accuracy for the same. The predicted values are geographically plotted.

4.10.6 KNN ALGORITHM

This module is used to design a K-Nearest Neighbour algorithm which is a semi-supervised model, get the sample data, training data from the database and finally predict the group it fit perfectly. The class values are helpful to get the first level of prediction values.

CHAPTER 5

RESULTS AND DISCUSSION

5.1 NEURAL NETWORKS VERSUS CONVENTIONAL COMPUTERS

Neural networks take a unique approach to hassle fixing than that of conventional computers. Conventional computers use an algorithmic approach i.e. The laptop follows a set of instructions on the way to solve a hassle. Solve the trouble. That restricts the trouble fixing functionality of conventional computers to issues that we already understand and recognize how to clear up. However computer systems would be so much more except the unique steps that the computer needs to observe are recognised the pc can't be beneficial if they may do things that we don't precisely recognise a way to do.

Neural networks procedure statistics in a comparable way the human mind does. The community is composed of a large range of quite interconnected processing elements (neurons) running in parallel to clear up a particular problem. Neural networks analyze by using instance. They can't be programmed to perform a specific venture. The examples have to be decided on cautiously in any other case beneficial time is wasted or even worse the community might be functioning incorrectly. The drawback is that because the community finds out how to resolve the trouble with the aid of itself, its operation may be unpredictable.

Then again, conventional computers use a cognitive approach to trouble fixing; the manner the hassle is to be solved need to be recognized and stated in small unambiguous commands. Those instructions are then transformed to a high stage language software and then into machine code that the pc can understand. These machines are completely predictable; if whatever is going wrong is because of a software program or hardware fault.

Neural networks and conventional algorithmic computer systems aren't in opposition however complement each other. There are obligations are more suitable to an algorithmic method like arithmetic operations and duties which are greater desirable to neural networks. Even extra, a big quantity of tasks, require

structures that use a mixture of the 2 tactics (typically a traditional laptop is used to supervise the neural community) that allows you to carry out at maximum efficiency.

5.2 Network layers

The most typical type of synthetic neural network includes 3 companies, or layers, of devices: a layer of "enter" units is hooked up to a layer of "hidden" units, which is related to a layer of "output" devices.

The activity of the enter gadgets represents the uncooked records this is fed into the network.

The pastime of every hidden unit is determined by the sports of the enter devices and the weights at the connections among the enter and the hidden devices.

The conduct of the output gadgets depends on the hobby of the hidden gadgets and the weights between the hidden and output units.

This easy kind of community is interesting because the hidden units are free to construct their very own representations of the input. The weights among the enter and hidden gadgets determine when every hidden unit is lively, and so by means of modifying those weights, a hidden unit can choose what it represents.

We also distinguish single-layer and multi-layer architectures. The unmarried-layer corporation, wherein all devices are connected to each other, constitutes the maximum general case and is of greater potential computational power than hierarchically dependent multi-layer corporations. In multi-layer networks, units are regularly numbered through layer, as opposed to following a worldwide numbering.

Normally neural networks are adjusted, or skilled, in order that a particular input results in a specific target output. This type of scenario is proven beneath. There, the community is adjusted, based on a comparison of the output and the goal, until the network output fits the goal. Commonly many such input/goal pairs are used, in this supervised learning, to educate a community.




Fig 5.2 block diagram of network layers

Batch schooling of a community proceeds by means of making weight and bias modifications based on an entire set (batch) of input vectors. Incremental training modifications the weights and biases of a network as wished after presentation of every man or woman input vector. Incremental education is sometimes referred to as “on line” or “adaptive” education.

Neural networks are composed of easy elements operating in parallel. These elements are stimulated through organic frightened structures. As in nature, the network feature is determined in large part through the connections between factors. We will train a neural community to perform a particular characteristic by adjusting the values of the connections (weights) among elements.

5.2.1 INITIALIZE THE NETWORK:

Permit’s begin with something easy, the introduction of a brand new network prepared for education.

Every neuron has a set of weights that want to be maintained. One weight for every input connection and an extra weight for the prejudice. We can want to shop additional homes for a neuron all through schooling, therefore we are able to use a dictionary to symbolize each neuron and store homes by using names which include ‘weights’ for the weights.

A network is organized into layers. The input layer is in reality just a row from our education dataset. The primary actual layer is the hidden layer. This is followed by way of the output layer that has one neuron for each magnificence fee.

We are able to organize layers as arrays of dictionaries and deal with the whole community as an array of layers.

It is right exercise to initialize the network weights to small random numbers. In this situation, can we use random numbers in the variety of zero to one.

Underneath is a feature named initialize community () that creates a new neural community prepared for education. It accepts three parameters, the number of inputs, the wide variety of neurons to have inside the hidden layer and the variety of outputs.

You may see that for the hidden layer we create n_hidden neurons and each neuron inside the hidden layer has n_inputs + 1 weights, one for each input column in a dataset and an additional one for the prejudice.

You may also see that the output layer that connects to the hidden layer has n_outputs neurons, every with n_hidden + 1 weight. Which means that each neuron within the output layer connects to (has a weight for) each neuron within the hidden layer.

5.3 TRAIN THE NETWORK

Once the community weights and biases are initialized, the network is ready for education. The multilayer feed forward network may be trained for function approximation (nonlinear regression) or pattern reputation. The education system requires a hard and fast of examples of right network conduct network inputs p and target outputs t. The system of training a neural community involves tuning the values of the weights and biases of the community to optimize network overall performance, as defined by using the network overall performance feature net. The default performance characteristic for feed ahead networks is mean square blunders mse—the common squared errors between the networks outputs and the target outputs t. It is described as follows:

$$F = mse = \frac{1}{N} \sum_{i=1}^N (e_i)^2 = \frac{1}{N} \sum_{i=1}^N (t_i - a_i)^2$$

There are different approaches wherein training can be carried out: incremental mode and batch mode. In incremental mode, the gradient is computed and the weights are up to date after each enter is carried out to the network. In batch mode, all of the inputs inside the training set are carried out to the network before the weights are updated. This topic describes batch mode schooling with the train command. Incremental schooling with the adapt command is discussed in Incremental schooling with adapt. For maximum troubles, whilst using the Neural network Toolbox™ software, batch education is notably quicker and produces smaller errors than incremental training. For education multilayer feed ahead networks, any popular numerical optimization set of rules may be used to optimize the performance characteristic, however there are a few key ones which have shown wonderful performance for neural community schooling. Those optimization strategies use both the gradient of the community overall performance with respect to the community weights, or the Jacobian of the network mistakes with recognize to the weights. The gradient and the Jacobian are calculated the usage of a way called the returned propagation set of rules, which entails acting computations backward through the community.

5.4 Training Algorithms:

As an instance of the way the training works, don't forget the only optimization algorithm — gradient descent. It updates the network weights and biases in the path wherein the performance characteristic decreases most hastily, the bad of the gradient. One generation of this set of rules can be written as

$$\mathbf{x}_{k+1} = \mathbf{x}_k - \alpha_k \mathbf{g}_k$$

Where $\mathbf{x}_k$ is a vector of cutting-edge weights and biases, $\mathbf{g}_k$ is the cutting-edge gradient, and α_k is the mastering fee. This equation is iterated till the network converges.

5.5 CODING

As soon as the design element of the gadget is finalized the system enters into the coding and checking out phase. The coding segment brings the actual machine into movement by way of converting the design of the gadget into the code in a given programming language. Consequently, an amazing coding fashion needs to be taken each time adjustments are required it easily screws into the system.

5.6 CODING STANDARDS

Coding requirements are suggestions to programming that makes a speciality of the physical structure and appearance of the program. They make the code easier to examine, apprehend and hold. This segment of the system definitely implements the blueprint evolved during the layout phase. The coding specification have to be in this type of way that any programmer have to be able to understand the code and may result in modifications on every occasion felt essential. Some of the same old had to acquire the above-referred to goals are as follows:

Program should be simple, clear and smooth to apprehend.

- Naming conventions
- Value conventions
- Script and comment procedure
- Message box format
- Exception and error handling

5.6.1 NAMING CONVENTIONS

Naming conventions of instructions, data member, member capabilities, methods and many others., must be self-descriptive. One must even get the which means and scope of the variable by its call. The conventions are followed for clean knowledge of the intended message by way of the consumer. So it is standard to comply with the conventions. Those conventions are as follows:

Class names

Class names are problem area equivalence and start with capital letter and have mixed instances.

Member function and facts Member name

Member function and records member call begins with a lowercase letter with every subsequent letters of the brand new words in uppercase and the relaxation of letters in lowercase.

5.6.2 VALUES CONVENTIONS

value conventions make sure values for variable at any point of time. This includes the following:

- proper default values for the variables.
- right validation of values in the subject.
- right documentation of flag values.

5.6.3 SCRIPT WRITING AND COMMENTING FASHIONABLE

Script writing is an art wherein indentation is utmost vital. Conditional and looping statements are to be properly aligned to facilitate clean knowledge. Feedback are included to decrease the wide variety of surprises that might occur whilst going through the code.

5.6.4 MESSAGE box format

whilst some thing must be triggered to the consumer, he ought to be able to understand it properly. To achieve this, a specific layout has been followed in displaying messages to the user. They are as follows:

- □X – person has executed unlawful operation.
- □! – information to the consumer.

5.7 OUTPUTS




Fig 5.7 input soil dataset




fig 5.8 preprocessing




Fig 5.9 train the neural network




Fig 5.10 pattern recognition neural network

CHAPTER 6

SUMMARY AND CONCLUSION

6.1 CONCLUSION

In this paper, the proposed system focused on implementing an efficient machine learning algorithm which is nothing but the combination of more than one ML algorithm to form a hybrid network in deep learning. Deep Hybrid learning model classify the various nutrients present in the tamilnadu location, district wise and formulates a pictorial model of Location classification too. Hybrid combination of KNN and SVM Model is developed. K-nearest neighbor is the semi-supervised model and it will be more effective in parameter level classification. The proposed model is apt for measuring the accuracy of hybrid model and readings will be tabulated.

REFERENCE

- [1] T. C. Wang, G. Qu, J. Li, and D. Liang, "Evaluation and optimization of multi-channel pulsed discharge plasma system for soil remediation," *Vacuum*, vol. 103, pp. 72–77, May 2014.
- [2] H. Zhang, D. Ma, R. Qiu, Y. Tang, and C. Du, "Non-thermal plasma technology for organic contaminated soil remediation: A review," *Chem. Eng. J.*, vol. 313, pp. 157–170, Apr. 2017.
- [3] H. Ölmez and M. Y. Akbas, "Optimization of ozone treatment of freshcut green leaf lettuce," *J. Food Eng.*, vol. 90, no. 4, pp. 487–494, 2009.
- [4] N. Ahsan, Y. Nanjo, H. Sawada, Y. Kohno, and S. Komatsu, "Ozone stress-induced proteomic changes in leaf total soluble and chloroplast proteins of soybean reveal that carbon allocation is involved in adaptation in the early developmental stage," *Proteomics*, vol. 10, no. 14, pp. 2605–2619, 2010.
- [5] C. J. Saitanis and M. G. Karandinos, "Effects of ozone on tobacco (*Nicotiana tabacum* L.) varieties," *J. Agronomy Crop Sci.*, vol. 188, no. 1, pp. 51–58, 2002.
- [6] S. S. Zambre, K. V. Venkatesh, and N. G. Shah, "Tomato redness for assessing ozone treatment to extend the shelf life," *J. Food Eng.*, vol. 96, no. 3, pp. 463–468, 2010.
- [7] L. Rodoni, N. Casadei, A. Concellón, A. R. C. Alicia, and A. R. Vicente, "Effect of short-term ozone treatments on tomato (*Solanum Lycopersicum* L.) fruit quality and cell wall degradation," *J. Agrc. Food Chem.*, vol. 58, no. 1, pp. 594–599, 2010.
- [8] X. C. Cui et al., "Arbuscular mycorrhizal fungi alleviate ozone stress on nitrogen nutrition of field wheat," *J. Agricult. Sci. Technol.*, vol. 15, no. 5, pp. 1043–1052, 2013.
- [9] N. Msayleb and S. Ibrahim, "Treatment of nematodes with ozone gas: A sustainable alternative to nematicides," *Phys. Procedia*, vol. 21, pp. 187–192, 2011.
- [10] M. Gómez-Alvarez, T. Poznyak, E. Ríos-Leal, and C. S. Sánchez, "Anthracene decomposition in soils by conventional ozonation," *J. Environ. Manage.*, vol. 113, pp. 545–551, Dec. 2012.
- [11] X. Bao, Q. Li, J. Hua, T. Zhao, and W. Liang, "Interactive effects of elevated ozone and UV-B radiation on soil nematode diversity," *Ecotoxicology*, vol. 23, no. 1, pp. 11–20, 2014.

- [12] J. Wang, X. Zhang, and G. Li, "Effects of ozonation on soil organic matter of contaminated soil containing residual oil," *J. Solids Sediments*, vol. 12, no. 2, pp. 117–127, 2012.
- [13] C. R. Yendrek, C. P. Leisner, and E. A. Ainsworth, "Chronic ozone exacerbates the reduction in photosynthesis and acceleration of senescence caused by limited N availability in *Nicotiana sylvestris*," *Global Change Biol.*, vol. 19, no. 10, pp. 3155–3166,
- [14] J. Rodriguez, A. García, T. Poznyak, and I. Chairez, "Phenanthrene degradation in soil by ozonation: Effect of morphological and physicochemical properties," *Chemosphere*, vol. 169, pp. 53–61,
- [15] H. Ikeura, F. Kobayashi, and M. Tamaki, "Removal of residual pesticide, fenitrothion, in vegetables by using ozone microbubbles generated by different methods," *J. Food Eng.*, vol. 103, no. 3, pp. 345–349, 2011.
- [16] H. Ikeura, F. Kobayashi, and M. Tamaki, "Removal of residual pesticides in vegetables using ozone microbubbles," *J. Hazardous Mater.*, vol. 186, no. 1, pp. 956–959, 2011.
- [17] Y.-C. Lin, H.-C. Juan, and Y.-C. Cheng, "Ozone exposure in the culture medium inhibits enterovirus 71 virus replication and modulates cytokine production in rhabdomyosarcoma cells," *Antiviral Res.*, vol. 76, no. 3, pp. 241–251, 2007.
- [18] K. R. Wigginton and T. Kohn, "Virus disinfection mechanisms: The role of virus composition, structure, and function," *Current Opinion Virol.*, vol. 2, no. 1, pp. 84–89, 2012.
- [19] A. A. Isikber and C. G. Athanassiou, "The use of ozone gas for the control of insects and micro-organisms in stored products," *J. Stored Products Res.*, vol. 64, pp. 139–145, Oct. 2015.
- [20] M. Nur, E. Kusdiyantini, W. Wuryanti, T. A. Winarni, S. A. Widyanto, and H. Muharam, "Development of ozone technology rice storage systems (OTRISS) for quality improvement of rice production," *J. Phys., Conf. Ser.*, vol. 622, no. 1, p. 012029, 2015.
- [21] F. Mendez, D. E. Maier, L. J. Mason, and C. P. Woloshuk, "Penetration of ozone into columns of stored grains and effects on chemical composition and processing performance," *J. Stored Products Res.*, vol. 39, no. 1, pp. 33–44, 2003.
- [22] M. X. McDonough, L. J. Mason, and C. P. Woloshuk, "Susceptibility of stored product insects to high concentrations of ozone at different exposure intervals," *J. Stored Products Res.*, vol. 47, no. 4, pp. 306–310, 2011.

- [23] F. Mitsugi, T. Abiru, T. Ikegami, K. Ebihara, S.-I. Aoqui, and K. Nagahama, "Influence of ozone generated by surface barrier discharge on nematode and plant growth," *IEEE Trans. Plasma Sci.*, vol. 44, no. 12, pp. 3071–3076, Dec. 2016.
- [24] F. Mitsugi, T. Abiru, T. Ikegami, K. Ebihara, and K. Nagahama, "Treatment of nematode in soil using surface barrier discharge ozone generator," *IEEE Trans. Plasma Sci.*, vol. 45, no. 12, pp. 3076–3081, Dec. 2017.
- [25] T. Nagatomo, T. Abiru, F. Mitsugi, K. Ebihara, and K. Nagahama, "Study on ozone treatment of soil for agricultural application of surface dielectric barrier discharge," *Jpn. J. Appl. Phys.*, vol. 55, no. 1S, p. 01AB06, 2015.
- [26] F. Mitsugi et al., "Properties of soil treated with ozone generated by surface discharge," *IEEE Trans. Plasma Sci.*, vol. 42, no. 12, pp. 3706–3711, Dec. 2014.
- [27] K. Ebihara et al., "Recent development of ozone treatment for agricultural soil sterilization and biomedical prevention," *Przegląd Elektrotechniczny*, vol. 88, no. 6, pp. 92–94, 2012.
- [28] T. Nagatomo, K. Takigawa, T. Sakai, F. Mitsugi, T. Ikegami, and K. Ebihara, "Fundamental study on ozone treatment of soil for agricultural application," in *Proc. 2nd IIAI Int. Conf. Adv. Appl. Inform.*, Aug./Sep. 2013, pp. 121–124.
- [29] F. Mitsugi, K. Ebihara, N. Horibe, S.-I. Aoqui, and K. Nagahama, "Practical soil treatment in a greenhouse using surface barrier discharge ozone generator," *IEEE Trans. Plasma Sci.*, vol. 45, no. 12, pp. 3082–3088, Dec. 2017.