

Journal Pre-proof

Ocean acidification affects biological activities of seaweeds: A case study of *Sargassum vulgare* from Ischia volcanic CO₂ vents

Amit Kumar, Maria Cristina Buia, Anna Palumbo, Mohamed Mohany, Mohammed A.M. Wadaan, Wael N. Hozzein, Gerrit T.S. Beemster, Hamada AbdElgawad

PII: S0269-7491(19)33161-6

DOI: <https://doi.org/10.1016/j.envpol.2019.113765>

Reference: ENPO 113765

To appear in: *Environmental Pollution*

Received Date: 17 June 2019

Revised Date: 6 December 2019


Accepted Date: 7 December 2019

Please cite this article as: Kumar, A., Buia, M.C., Palumbo, A., Mohany, M., Wadaan, M.A.M., Hozzein, W.N., Beemster, G.T.S., AbdElgawad, H., Ocean acidification affects biological activities of seaweeds: A case study of *Sargassum vulgare* from Ischia volcanic CO₂ vents, *Environmental Pollution* (2020), doi: <https://doi.org/10.1016/j.envpol.2019.113765>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier Ltd.





Ocean acidification affects biological activities of seaweeds: A case study of *Sargassum vulgare* from Ischia volcanic CO₂ vents

Amit Kumar^{a,b}, Maria Cristina Buia^c, Anna Palumbo^d, Mohamed Mohany^e, Mohammed A.M. Wadaan^f, Wael N. Hozzein^{f,g}, Gerrit T.S. Beemster^h, Hamada AbdElgawad^g

^aCentre for Climate Change Studies, Sathyabama Institute of Science and Technology, Jeppiaar Nagar, Rajiv Gandhi Salai, Chennai, India

^bSathyabama Marine Research Station, 123 Sallimalai Street, Rameswaram, India

^cCenter of Villa Dohrn Ischia – Benthic Ecology, Department of Integrative Marine Ecology, Stazione Zoologica Anton Dohrn, P.ta S. Pietro, Ischia, Naples, Italy

^dDepartment of Biology and Evolution of Marine Organisms, Stazione Zoologica Anton Dohrn, Villa Comunale, Naples, Italy

^eDepartment of Pharmacology and Toxicology, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia

^fBioproducts Research Chair, Zoology Department, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

^gBotany and Microbiology Department, Faculty of Science, Beni-Suef University, Beni-Suef, Egypt

^hIntegrated Molecular Plant Physiology Research Group (IMPRES) Department of Biology, Groenenborgerlaan 171, University of Antwerp, Antwerp, Belgium

Corresponding Author: Amit Kumar, E-mail: amitkumar.cccs@sathyabama.ac.in

Abstract

We utilized volcanic CO₂ vents at Castello Aragonese off Ischia Island as a natural laboratory to investigate the effect of lowered pH/elevated CO₂ on the bioactivities of extracts from fleshy brown algae *Sargassum vulgare* C. Agardh. We analysed the carbohydrate levels, antioxidant capacity, antibacterial, antifungal, antiprotozoal, anticancer properties and antimutagenic potential of the algae growing at the acidified site (pH ~ 6.7) and those of algae growing at the nearby control site Lacco Ameno (pH~8.1). The results of the present study show that the levels of polysaccharides fucoidan and alginate were higher in the algal population at acidified site. In general, extracts for the algal population from the acidified site showed a higher antioxidant capacity, antilipidperoxidation, antibacterial, antifungal, antiprotozoal, anticancer activities and antimutagenic potential compared to the control population. The increased bioactivity in acidified population could be due to elevated levels of bioactive compounds of algae and/or associated microbial communities. In this snapshot study, we performed bioactivity assays but did not characterize the chemistry and source of presumptive bioactive compounds. Nevertheless, the observed improvement in the medicinal properties of *S. vulgare* in the acidified oceans provides a promising basis for future marine drug discovery.

Capsule: Elevated DIC improves bioactivity in fleshy brown macroalgae at natural volcanic CO₂ off Ischia Island, Italy.

Keywords: Climate change; Bioactivity; Marine algae; Natural acidification; Marine drugs

1. Introduction

Volcanic CO₂ vents off Ischia Island in the Mediterranean Sea offer a unique setting to investigate the effects of ocean acidification (OA) on marine organisms (Foo et al., 2018). These vents are in particular useful to investigate benthic biota with less mobility and limited dispersal of propagules or larvae and that have occupied these areas for years (Foo et al., 2018). Venting activities have created a pH gradient along the rocky shores. Decreasing pH has caused a reduction in calcifying species (Hall-Spencer et al., 2018). At the most acidified site (pH ~6.7), calcareous algae has disappeared and species richness is decreased by 72% in comparison to control sites higher in the pH gradient (pH~8.1). The acidified site is dominated by fleshy seaweeds, in particular the brown algae *Sargassum vulgare* (Hall-Spencer et al., 2008; Porzio et al., 2011; Gao et al., 2019). This natural CO₂ gradient provides a unique set up to investigate how lowered pH/elevated dissolved inorganic carbon (DIC) may affect bioactivity of *S. vulgare* and associated microorganisms.

Theoretically fleshy seaweed should benefit from elevated carbon dioxide (CO₂) and bicarbonate (HCO₃⁻) as it can be used for photosynthesis (Koch et al., 2013; Mackey, 2015). However, growing evidence from the literature suggests that the response of seaweeds to ocean acidification depend on which DIC source it uses as well as mechanisms of DIC acquisition (Koch et al., 2013). There are three major DIC in seawater: CO₂ (~1%), carbonate (CO₃²⁻ ~8%) and bicarbonate (HCO₃⁻ ~91%), of which only CO₂ and HCO₃⁻ can be used for photosynthesis (Roleda and Hurd, 2012). Though CO₂ can diffuse inside the cell, its low concentration and low diffusion rate in seawater limits its availability for photosynthesis. Hence, almost 95% of marine macrophytes have developed efficient carbon concentrating mechanisms (CCM) i.e. uptake of HCO₃⁻ from the surrounding water inside the cell for photosynthesis (Koch et al., 2013). CCMs includes the active uptake of HCO₃⁻ involving P-type H⁺ ATPase pump, direct uptake of HCO₃⁻ through plasmalemma-located anion exchange (AE) proteins, and involvement of external and internal carbon anhydrase (CA)

enzyme activity (Fernández et al., 2014). External CA dehydrates HCO_3^- into CO_2 , which diffuses inside the cell where internal CA interconvert CO_2 to HCO_3^- to maintain internal pH and avoid CO_2 leakage (Fernández et al., 2018). Furthermore, chloroplast CA ensures the supply of CO_2 to the active site of RuBisCO for carbon fixation (Fernández et al., 2018). Thus, seaweeds lacking CCMs are likely to benefit from additional aqueous CO_2 (Kübler et al., 1999), but even seaweeds with CCMs may benefit from excess aqueous CO_2 if they can simultaneously use CO_2 and HCO_3^- under ambient and ocean acidification conditions to reduce the energy costs (Fernández et al., 2015; Wu et al., 2008). In contrast to photosynthesis, other physiological processes including ion homeostasis, respiration, nutrient uptake, and enzymatic activities are inhibited by ocean acidification conditions (Fernández et al., 2015; Gutow et al., 2014; Hofmann et al., 2013; Roleda et al., 2012).

Around the world, seaweeds are utilized as food and medicines since prehistoric times (Mouritsen et al., 2018). They are rich sources of proteins, vitamins, lipids, sugars, dietary fibre, and nutrients essential for human nutrition (Perumal et al., 2019). They are also known for their importance in pharmacology, being a rich source of bioactive compounds against bacteria, fungi, virus, protozoa, and various cancers (Hao et al., 2019; Smit et al., 2004). Because of these benefits, seaweeds are gradually gaining importance as healthy food (Cherry et al., 2019). The algal genus *Sargassum* (Ochrophyta, Phaeophyceae), in particular has anticancer, antibacterial, antifungal, anti-viral, anti-inflammatory, anticoagulant, antioxidant, hypoglycaemic, hypolipidemic, antimelanogenic, antibone loss, hepatoprotective and neuroprotective activities (Ghanian et al., 2017; Liu et al., 2012; Pramesti et al., 2017; Torres et al., 2014). Bioactivities of the seaweeds are attributed to primary and secondary metabolites e.g. meroterpenoids, phlorotannins, polysaccharides, phytosterols, polyunsaturated fatty acids and glycolipids (Dore et al., 2013; Hao et al., 2019; Kolsi et al., 2017; Liu et al., 2012; Sousa et al., 2008; Torres et al., 2007).

The effects of global climate change mainly through ocean acidification (OA) and warming have the potential to alter the carbon physiology of seaweeds and levels of metabolites, including bioactive compounds and nutritional value (Yousefzadi & Miejalili, 2017). Increased CO₂ concentration increase level of sugars, essential amino acids, phenolic, and vitamins, which in-turn improves total antioxidant capacity, and bioactivity against several microbes and cancer cells (Al Jaouni et al., 2018; Ghasemzadeh, 2011; Idso et al., 2000; Jaafar et al., 2012; Saleh et al., 2018). So far only a handful of studies have reported nutritional changes in seaweed under acidified conditions. The level of proteins and lipids increased in the green algae *U. rigida* in response to ocean acidification (Gao et al., 2017). OA also increased the colour and flavour of red algae *Pyropia yezoensis* by increasing the levels of pigments, and amino acids (Gao et al., 2019). Several seaweeds exposed to OA conditions showed varying responses on the levels of phenols and flavonoid compounds (Arnold et al., 2012; Betancor et al., 2014; Celis-Plá et al., 2015; Celis-Plá et al., 2017; Kumar et al., 2018). Moreover, the severity of the impact of ocean acidification on seaweeds depends on other environmental parameters including nutrients (Gao et al., 2018). It is therefore important to better understand how OA will impact the metabolite and bioactive properties of seaweeds.

In the present study, we investigated bioactive properties of *S. vulgare* growing naturally at the most acidified site of the volcanic CO₂ vents and control site off the Ischia Island, Italy. Previous studies have shown *S. vulgare* can utilize both CO₂ and HCO₃⁻ as a carbon source for photosynthesis and may therefore not be carbon limited (Kumar et al., 2017a). Nevertheless, this alga showed significant increase in photosynthetic performance (relative electron transport rate and saturating irradiance) upon short term exposure to elevated DIC (Kumar et al., 2017a). They exhibit high plasticity to elevated DIC/lowered pH conditions by activating molecular and physiological strategies to grow in the acidified water (Kumar et al., 2017a, Kumar et al., 2017b). Primary and secondary metabolites of *S. vulgare* were strongly

influenced, in particular the level of soluble carbohydrates were increased (Kumar et al., 2018; Porzio et al., 2017). Because soluble carbohydrates can act as precursors for bioactive metabolites, the current study is aimed to investigate the possibility that OA increase bioactive properties of *S. vulgare*. Therefore, we analysed the levels of polysaccharides fucoidan and alginate and antioxidant capacity, antibacterial, antifungal, antiprotozoal, anticancer properties and antimutagenic potential of the algae growing at the acidified site (pH ~ 6.7) and compared to those of algae growing at the nearby control site Lacco Ameno (pH~8.1). The present work represents a snapshot study where we conducted bioactivity assays but did not characterize the chemistry and source of presumptive bioactive compounds.

2. Materials and Methods

2.1 Sampling locations and seaweed collection

Sargassum vulgare (Class Phaeophyceae, Order Fucales, Family Sargassaceae) was collected from two different locations along Ischia Island, Italy: Castello Aragonese (40°43.87N, 013°57.78E) – the acidified site, and Lacco Ameno - the control site (40°45.35N, 013°53.13E, Fig. 1). Castello Aragonese site is characterized by acidified seawater due to continuous emission of CO₂ (90.1–95.3%) from underwater volcanic vents, for 2 millennia (Lombardi et al., 2011), with no detectable harmful sulfur gases. Though the venting activities are quite variable, on average pH values, where *S. vulgare* occurs are constantly around 6.72 (Porzio et al., 2011). Lacco Ameno almost 6 km northwest from Castello Aragonese, was used as a control site as algae are growing here at the same hydrodynamic and physical conditions but at current pH values (8.1). In both locations, *S. vulgare* grows at <1 m depth, in a sheltered bay with similar geographical exposure (Kumar et al., 2017b). Salinity fluctuates between 37 and 38.5 psu at both the sites, which is typically the case in Tyrrhenian coastal waters (d'Alcalà et al., 2004; Lorenti et al., 2005). Temperature varies between 14 - 28 °C during the year and shows no significant difference between the

sites (Kumar et al., 2017b). Photosynthetically active radiation (PAR) irradiances obtained by the PAR spherical sensor QSI-140 (Biospherical Instruments, San Diego, USA) as K_d coefficient from surface to 1 m depth was reported to be similar (0.18 and 0.20 at acidified and control site, respectively) (Lorenti & De Falco 2004). Therefore, we believe that depth, salinity, temperature, and light were not confounding factors. Carbon chemistry of acidified site has been assessed during the last 10 years (Calosi et al., 2013; Cigliano et al., 2010; Hall-Spencer et al., 2008; Kroeker et al., 2013; Kumar et al., 2017b; Ricevuto et al., 2014), and control site assessed during Apr-May 2013 and described by Ricevuto et al., (2015) is summarized in Table A.1 in Appendices. Total alkalinity was similar at both sites, while $p\text{CO}_2$, HCO_3^- , and DIC contents were higher at acidified site, and CO_3^{2-} was higher at the control site (Ricevuto et al., 2014). Nutrient levels at both sites varied, e.g. nitrate was similar in both sites, while phosphate was almost 40% lower and potassium was 27% higher in acidified waters than control site (Kumar et al., unpublished).

Seaweed thalli were collected at both sites on the same day (March, 2014 at midday) at the same time (11AM-12 PM) to avoid influence of unpredictable and difficult to measure environmental factors. A total of 9 individual thalli of similar frond length (8-10 cm) were handpicked along a 15 m coastal stretch to cover natural variability of the local population at each site. Upon collection, the algae were maintained in seawater from respective sites and transported to the laboratory within an hour. The algal tissues were washed with filtered sea water to remove visible epiphytes, flash frozen in liquid nitrogen and immediately stored at -80°C until further analysis. Determination of fucoidan, alginate and biological activity assessment, were performed on at least three independent thalli from both the locations, thus creating three biological replicates from each site.

2.2 Determination of Fucoidan and Alginate

Fucoidan in the algal tissue was determined according to Turan et al. (2017). Briefly, 50 mg of algal tissue was ground using a MagNALyser (Roche, Vilvoorde, Belgium) and 2

ml of 0.2 N HCl was added and the suspension was heated for 1 h at 70 °C with regular stirring followed by cooling so that particles were allowed to settle. The liquid was filtered, stored and then the retained particles were again extracted using 5 ml of 0.2N HCl. Finally, both liquids were combined together. 1ml of sample extract was mixed with 4.5 ml dilute H₂SO₄ in the ice-water bath for ~1 min followed by boiling for 10 min. Upon cooling to room temperature, 0.1 ml of Cysteine solution was added, mixed and incubated at room temperature for 30 minutes. The absorbance in the solution was observed at 396 and 427 nm for fucose content which was multiplied by a factor of 2 to obtain fucoidan content.

For alginate determination, the algal tissue (50 mg) was extracted in 2 ml 0.1 M sodium citrate and centrifuged for 30 min at 5000 rpm. To separate alginate, supernatant was treated with 1.5 M hydrochloric acid for 1 h. After precipitation, tubes were centrifuged and the precipitates were mixed with 500 µl citrate buffer, subsequently alkalinised with 500 µl of 0.8N NaOH, finally neutralized after 5 min by adding 60µl of 2.25N citrate. The colorimetric estimation was done by adding 20µl of 1, 9-dimethyl methylene blue (DMMB) in the homogenate, measured at 520 nm and 650 nm and calculated as described by Hallé et al. (1993).

2.3 Biological activity assessments

2.3.1 *In vitro* antioxidant activity

Total antioxidant capacity was determined utilizing ferric reducing antioxidant power (FRAP) and diphenylpicrylhydrazyl (DPPH) assays. 300 mg algal tissue was ground using a MagNALyser (Roche, Vilvoorde, Belgium) and mixed with 3 ml of 80% ice-cold ethanol. The homogenate was centrifuged for 30 min at 14000 rpm at 4 °C and the supernatant was collected. FRAP assay was performed according to Abdelgawad et al. (2015). Briefly, 20 µl of the ethanol extract was mixed with 180 µl of freshly prepared pre-warmed FRAP reagent (10 mM 2, 4, 6 Tris (2-ylidyl) s-triazine (TPTZ), 0.3 M acetate and 20 mM FeCl₃) and incubated for 30 min at 37 °C. Absorbance of the mixture was measured at 593 nm using a

microplate reader (Synergy Mx, Biotek Instruments Inc., Vermont, VT, USA). Trolox was used to prepare standard curve and using slope and intercept, antioxidant concentration was determined which was expressed as $\mu\text{MTrolox equivalent g}^{-1}$ frozen weight.

DPPH assay was performed according to Al Jaouni et al. (2015). Briefly, algal extract was prepared by grinding 300 mg of tissue in 3 ml of 80% ice-cold ethanol and mixed with 0.5 ml of DPPH solution (0.25 mM in 95% ethanol, followed by incubation at room temperature for 30 min. Thereafter, absorbance of the mixture was measured at 517 nm to calculate the inhibition percentage.

2.3.2 Antilipid peroxidation assay

The degree of lipid peroxidation was measured by the thiobarbituric acid reactive substance (TBARS) method, using egg yolk homogenate as lipid rich substrate (Ohkawa et al., 1979). Algal extract was prepared by grinding 100 mg algal tissue using a MagNALyser (Roche, Vilvoorde, Belgium) and mixed with 2 ml of 90% ethanol. Algal extract and egg homogenate (0.5ml of 10% v/v) were mixed with 15 mM ferrous sulphate (to induce lipid peroxidation), after incubation for 30 min, 1.5 ml of 10% TCA was added. Following that, the mixture was transferred to a tube containing 1.5 ml of 0.67% TBA and boiled for 30 min. The chromogen formed was measured at 535 nm.

2.3.3 Antimicrobial activities

Algal extract was prepared by mixing 100 mg of grinded tissue in 90% ethanol and then evaporated using centrifugal rotor evaporator. The antibacterial activity was determined using the disc diffusion method (7.5 mg of the algal ethanol extract per filter paper disc) on Muller Hinton agar (MHA inoculated with 100 μl of a suspension containing 10^8 CFU/ml of bacteria) (Selim et al., 2013). The plates were incubated at 35 °C for 24 h and the activity was expressed as the diameter of the inhibition zone in mm. Negative controls were prepared using the disc containing ethanol only.

2.3.4 Antiprotozoal activities

In vitro assessment of the antiprotozoal activity was performed according to Răz et al. (1997). *Trypanosoma cruzi* and *Leishmania donovani* were cultured in Minimum Essential Medium with additional supplements. Algal extract prepared in 90% ethanol and then dried in rotor evaporator (5 mg/ml) were used against *T. cruzi* and *L. donovani* and compared with control having no algal extract. The results were expressed as percentage of reduction in parasite load in treated and control conditions.

2.3.5 Antimutagenicity assay (Ames test)

An Ames assay (Maron and Ames, 1983) was performed to measure the antimutagenicity of β -D-glucogallin. Two strains of *Salmonella typhimurium* (TA98 and TA100) were used for the test. The assay was conducted in pre-incubation and co-incubation mode using 100 μ l of overnight grown bacterial cultures with the addition of algal extract and mutagenic agents in triplicates. The algal extract was prepared by grinding 100 mg tissue in ethanol/ water (1:1 v/v) followed by filtration using 0.45 μ m filter (Millipore, USA). Overnight bacterial culture, mutagen, algal extract and phosphate buffer were added to agar containing histidine/biotin (0.5 mM). As a positive control, Mutagen 4-nitro-o-phenylenediamine (NPD) and NaN_3 were used for TA98 and TA100 and ethanol/water (1:1, v/v) was used as negative control instead of algal extract. The mixture was poured onto minimal glucose plates. Histidine independent revertants colonies and viable cells were scored on plates after incubation at 37°C for 48 or 72 h. After incubation for 48 hrs at 37 °C, the number of revertants per plate was counted. The inhibition percent of mutageneic activity was calculated by observing the difference in the number of histidine revertants induced by mutagen alone, mutagen in the presence of the extract and extract alone.

2.3.6 Anticancer activities

The anticancer potential of the algal extract was assessed against 4 human cancer cell lines: hepatocellular carcinoma (HepG2), colon carcinoma (Colo205), embryonic kidney adenocarcinoma (293), and urinary bladder carcinoma (T24P). Algal extract prepared in 90%

ethanol and then dried in rotor evaporator. To assess the activity of the algal extract against these cell lines, cells were grown in Dulbecco's Modified Eagle Medium (DMEM, fetal calf serum, Na-pyruvate, 1mg/0.1 ml streptomycin, 1U/ 0.1ml penicillin) in the presence and absence of algal extract (2 mg/ml) at 37°C and 5% CO₂. Then, cells were harvested through trypsinization (Saleh et al., 2018). Cell viability was determined using a Cell Titer-Blue reagent as described by Solowey et al. (2014) and the results were expressed as percentage dead cells.

2.4 Statistical Analysis

Statistical analyses were performed using SPSS (v21 SPSS Inc, Chicago, IL, USA). The data were checked for normality by visual inspection and homogeneity of variance by Levene's test before applying independent sample t-test on the mean data values to determine significant differences between control and acidified site samples.

3. Results

To investigate the effect of ocean acidification on the metabolic composition of *S. vulgare*, we analysed the thalli growing naturally under acidified conditions and control conditions. The concentration of fucoidan was significantly higher in algae grown at the acidified site (46.71 ± 1.41 mg/FrozendryW) compared to those from the control site (34.96 ± 0.40 mg/gFrozendryW). Similarly, alginate content was also found to be higher in algae at the acidified site (93.3 ± 2.98 mg/gFrozendryW) than in those growing at the control site (74.1 ± 2.02 , Fig. 2).

When compared to algae from control site, algae from the acidified site showed improved diphenylpicrylhydrazyl (DPPH) antioxidant scavenging activities ($25.40 \pm 0.23\%$ vs $15.51 \pm 0.11\%$, $P < 0.0001$), however, no significant differences were observed in ferric reducing antioxidant power (FRAP) antioxidant and anti-lipid peroxidation activities (Fig. 3A, B, C).

Under elevated dissolved CO₂ conditions at the volcanic vents of Ischia, *S. vulgare* showed a general increase in the bioactivity against bacteria, fungi, protozoa, and cancer cells. The significant increase in antibacterial activity was recorded for *Pseudomonas aeruginosa* (p=0.00013), followed by *Klebsiella pneumonia* (p=0.008), *Escherichia coli* (p=0.003), *Proteus vulgaris* (p=0.002), *Serratia marcescens* (p=0.04), *Streptococcus sp.* (p=0.002). Only for *S. aureus*, no significant change (p=0.053) in the activity was observed (Fig. 4). The antifungal activities were significantly higher in the acidified samples for *Aspergillus parasiticus* (p=0.0003), and *Fusarium oxysporum* (p=0.008), however, no difference in activity was observed against *Candida albicans* (p=0.53), *Penicillium expansum* (p=0.07), and *Trichoderma harzianum* (0.559). The antiprotozoal activities showed that algal extracts from the acidified site exhibited significant lower values for IC₅₀ than from control sites against *Trypanosoma cruzi* (p=0.001), and *Leishmania donovani* (p=0.0024) (Fig. 5). The anticancer activities against all four cancer cell lines, HepG2 (p=0.0001), Colo205 (p=0.0001), T24P (p=0.002), and 293 cells (0.0002) tested were more pronounced in the algal extracts from the acidified site (Fig. 6). In the antimutagenicity assays, performed with TA98 and TA100, the algal extract from the acidified site showed higher percent inhibition of revertants in TA 98 (p= 0.001) and TA100 (p=0.03) than algal extracts from the control site (Fig. 7). The output of t-test is given in Table A. 2 as appendices.

4. Discussion

In this study, we tested the hypothesis that increased carbohydrate availability in *S. vulgare* exposed to elevated DIC at Ischia CO₂ vents can increase the bioactivity against microbial pathogens and cancer cells. Our findings supporting this hypothesis includes the elevated levels of polysaccharides fucoidan and alginate and increased antioxidant capacity, antilipidperoxidation, antibacterial, antifungal, antiprotozoal, anticancer activities and antimutagenic potential in algal homogenate from the population at acidified site compared to the control population.

4.1 Elevated dissolved CO₂ increases the content of polysaccharides

Anionic polysaccharide alginates and fucoidan measured in the present study are major constituents of the brown algal cell wall. The elevated level of these sugars could be due to increased photosynthesis in elevated DIC conditions in the short term despite the presence of CCM in *S. vulgare* (Kumar et al., 2017a). Since, *S. vulgare* is a mixed CO₂ and HCO₃⁻ user, elevated CO₂ might have down regulated direct or indirect HCO₃⁻ acquisition and the energy saved by bypassing anhydrase activity could be used for enhanced growth (Fernández et al., 2015) in the volcanic CO₂ seeps. As such, there is no clear mechanistic understanding regarding CO₂ concentrations and cell wall sugars, however, it is reported that higher CO₂ concentrations can increase the activity of enzymes involved in the synthesis of cell wall uronic acid, thus increasing synthesis of cell wall polysaccharide (Cheng et al., 2015). In our previous molecular study, we found that genes encoding for the enzymes related to the formation and structures of cell walls and carbon storage were expressed at higher levels in *S. vulgare* at elevated DIC conditions than in control conditions (Kumar et al., 2017b). In other seaweeds, high carbohydrate concentration and improved photosynthetic rates were also reported under elevated CO₂ (Figueroa et al., 2014; Suárez-Álvarez et al., 2012).

4.2 Elevated dissolved CO₂ increases bioactivity

The bioactivity of seaweeds is attributed to its primary and secondary metabolites including fucoxanthin, alginates, laminarins, fucans, cellulose as well as unsaturated fatty acids and essential amino acids (Dore et al., 2013; Kolsi et al., 2017; Liu et al., 2012; Sousa et al., 2008; Torres et al., 2007). Acting as a substrate for photosynthesis, elevated CO₂ (up to certain physiological limit) has the potential to enhance carbon assimilation, thus improving the accumulation of non-structural carbohydrates and their breakdown through dark reaction (Leahey et al., 2009). This might provide increased levels of precursors and energy for the synthesis of different bioactive secondary metabolites (Kumar et al., 2018; Saleh et al., 2018).

In the current study, elevated CO₂ resulted in increased levels of polysaccharide fucoidans and alginates which are known to have antimicrobial and anticancer bioactivity (Costa et al., 2011; Jun et al., 2018; Sousa et al., 2008; Wu et al., 2016). In *S. vulgare* increased levels of fucoidans and alginates has been related to improved antimicrobial and anticancer activity (Wu et al., 2016) as well as antiprotozoal activity (Armeli Minicante et al., 2016). *S. vulgare* at acidified site has increased the level of photosynthetic pigments (Kumar et al., 2017a), which may be linked with improved antimutagenicity as pigments either suppress the metabolic activation of mutagens or inhibit activated mutagens (Osuna-Ruiz et al., 2016; Hayatsu et al., 1993).

We demonstrated that the antioxidant capacity of algal extracts varied depending on the assay used. With DPPH radical scavenging method, the antioxidant potential of algae of the acidified site was higher compared to control population, whereas no difference was observed with FRAP assay. This is not surprising considering that the two assays measured different parameters. Whereas the FRAP method is based on the ability of antioxidant to reduce ferric ion complex, the DPPH method measures the reduction of an organic radical by the capability of the antioxidant to donate a hydrogen (Csepregi et al., 2016). Antioxidant capacity in seaweeds is mostly attributed to polyphenols (Machu et al., 2015). Even though lowered levels of polyphenols and flavonoid were found in the algae at the acidified site in comparison to control site (Kumar et al., 2017a; Kumar et al., 2018), it is interesting that DPPH radical scavenging is higher in these algae. *Sargassum* contains more than 280 metabolic compounds including, pigments, primary and secondary metabolites with bioactive properties (Liu et al., 2012). Therefore, our result suggests that other bioactive molecules which are not yet identified may be contributing towards antioxidant activity of *S. vulgare* (Kumar et al., 2018). Another reason for dissimilar results between metabolites and bioactivity may be due to different sampling season. Moreover, temporal and spatial variation has been reported in biochemical composition and metabolite contents of seaweeds (Graiff et

al., 2015; Pavia et al., 2018; Roleda et al., 2019; Véliz et al., 2018). Seasonal variation in bioactivity of the genus *Sargassum* has been reported (Praioboon et al., 2015). It is also important to note that recent studies on seaweed associated microorganisms (holobiont) indicated that at least part of bioactivity observed in the present study could be from associated microbial communities (Singh and Reddy, 2014). Also, associated microbial communities are reported to be site and tissue specific in the seaweeds (Serebryakova et al., 2018; Campbell et al., 2011), and microbial community structure changes under acidified conditions (Aires et al., 2018). Hence, the involvement of microbes in these bioactivities cannot be ruled out in the present study. In future, manipulative laboratory experiments will be conducted by bringing samples from both sites and expose to ocean acidification to eliminate the role of associated microorganisms and to ensure the observed bioactivity is due to elevated DIC or lowered pH or any other factors.

In conclusion, the present work on *S. vulgare* expands our understanding of the bioactivity derived from seaweeds under a future climate scenario. As an outcome, this preliminary study can be seen as a positive sign for marine drug discovery considering the improvement in the medicinal properties of *S. vulgare* or associated microbial communities in the acidified oceans. It would be interesting to conduct a larger scale metabolite profiling analysis to understand which compounds contribute towards algal bioactivity, source of bioactive compounds (whether seaweed directly or associated microbes) and to investigate biochemical pathways leading to varying metabolite levels that are activated by ocean acidification.

Acknowledgment

This work was co-funded by SZN and the Flagship RITMARE – The Italian Research for the Sea – coordinated by the Italian National Research Council and by the Italian Ministry of Education, University and Research within the National Research Program 2012–2015. The authors extend their appreciation to the Deanship of Scientific Research at King Saud

University for funding this work through the research group no. RGP-205. HAE was supported by a postdoctoral fellowship from the Flemish Science Foundation (FWO, 12U8918N). We thank Captain V. Rando for his assistance in fieldwork.

References

- AbdElgawad, H., Farfan-Vignolo, E. R., Vos, D. d., Asard, H., 2015. Elevated CO₂ mitigates drought and temperature-induced oxidative stress differently in grasses and legumes. *Plant Sci.* 231, 1-10. <https://doi.org/10.1016/j.plantsci.2014.11.001>.
- Aires, T., Serebryakova, A., Viard, F., Serrão, E.A., Engelen, A.H., 2018. Acidification increases abundances of Vibrionales and Planctomycetia associated to a seaweed-grazer system: potential consequences for disease and prey digestion efficiency. *Peer J.* 6, e4377. <https://doi.org/10.7717/peerj.4377>.
- Al Jaouni, S., Saleh, A.M., Wadaan, M. A. M., Hozzein, W.N., Selim, S., AbdElgawad, H., 2018. Elevated CO₂ induces a global metabolic change in basil (*Ocimum basilicum* L.) and peppermint (*Mentha piperita* L.) and improves their biological activity. *J. Plant Physiol.* 224 121-131. <https://doi.org/10.1016/j.jplph.2018.03.016>.
- Armeli Minicante, S., Michelet, S., Bruno, F., Castelli, G., Vitale, F., Sfriso, A., Morabito, M., Genovese, G., 2016. Bioactivity of phycocolloids against the mediterranean protozoan *Leishmania infantum*: An Inceptive Study. *Sustainability.* 8, 1131. <https://doi.org/10.3390/su8111131>.
- Arnold, T., Mealey, C., Leahey, H., Miller, A.W., Hall-Spencer, J.M., Milazzo, M., Maers, K., 2012. Ocean acidification and the loss of phenolic substances in marine plants. *PLoS One* 7, e35107. <https://doi.org/10.1371/journal.pone.0035107>.
- Betancor, S., Tuya, F., Gil-Díaz, T., Figueroa, F.L., Haroun, R., 2014. Effects of a submarine eruption on the performance of two brown seaweeds. *J. Sea Res.* 87, 68–78. <https://doi.org/10.1016/j.seares.2013.09.008>.

- 414 Calosi, P, Rastrick S.P.S., Lombardi, C., de Guzman, H.J., Davidson, L., Jahnke, M.,
 415 Giangrande, A., Hardege, J.D., Schulze, A., Spicer, J.I., Gambi, M.C., 2013. Adaptation
 416 and acclimatization to ocean acidification in marine ectotherms: an in situ transplant
 417 experiment with polychaetes at a shallow CO₂ vent system. Philos. T R Soc. B. 368,
 418 20120444-20120444. <https://doi.org/10.1098/rstb.2012.0444>.
- 419 Campbell, A.H., Harder, T., Nielsen, S., Kjelleberg, S., Steinberg, P.D., 2011. Climate
 420 change and disease: bleaching of a chemically defended seaweed. Glob. Change Biol.
 421 17, 2958-2970. <https://doi.org/10.1111/j.1365-2486.2011.02456.x>.
- 422 Celis-Plá, P.S., Hall-Spencer, J.M., Horta, P.A., Milazzo, M., Korbee, N., Cornwall, C.E.,
 423 Figueroa, F.L., 2015. Macroalgal responses to ocean acidification depend on nutrient
 424 and light levels. Front. Mar. Sci. 2, 26. <https://doi.org/10.3389/fmars.2015.00026>.
- 425 Celis-Plá, P.S., Martínez, B., Korbee, N., Hall-Spencer, J.M., Figueroa, F.L., 2017.
 426 Ecophysiological responses to elevated CO₂ and temperature in *Cystoseira*
 427 *tamariscifolia* (Phaeophyceae). Clim. Change 142, 67–81. [https://doi.org/10.1007/s10584-](https://doi.org/10.1007/s10584-017-1943-y)
 428 017-1943-y.
- 429 Cheng, Y. , Labavitch, J., VanderGheynst, J., 2015. Elevated CO₂ concentration impacts cell
 430 wall polysaccharide composition of green microalgae of the genus *Chlorella*. Lett.
 431 Appl. Microbiol., 60, 1-7. <https://doi.org/10.1111/lam.12320>.
- 432 Cherry, P., O'Hara, C., Magee, P.J., McSorley, E.M., Allsopp, P.J., 2019. Risks and benefits
 433 of consuming edible seaweeds. Nutr. Rev. 77, 307-329.
 434 <https://doi.org/10.1093/nutrit/nuy066>.
- 435 Cigliano, M., Gambi, M., Rodolfo-Metalpa, R., Patti, F., Hall-Spencer, J., 2010. Effects of
 436 ocean acidification on invertebrate settlement at volcanic CO₂ vents. Mar. Biol. 157,
 437 2489-2502. <https://doi.org/10.1007/s00227-010-1513-6>.
- 438 Costa, L.S., Fidelis, G.P., Telles, C.B.S., Dantas-Santos, N., Camara, R.B.G., Cordeiro, S.L.,
 439 Costa, M.S.S.P., Almeida-Lima, J., Melo-Silveira, R.F., Oliveira, R.M., Albuquerque, I.

- 440 R.L., Andrade, G.P.V., Rocha, H. A. O., 2011. Antioxidant and antiproliferative
441 activities of heterofucans from the seaweed *Sargassum filipendula*. Mar. Drugs. 9, 952-
442 966. [https://doi: 10.3390/md9060952](https://doi.org/10.3390/md9060952).
- 443 Csepregi, K., Neugart, S., Schreiner, M., Hideg, É., 2016. Comparative Evaluation of Total
444 Antioxidant Capacities of Plant Polyphenols. Molecules 21, 208.
445 <https://doi.org/10.3390/molecules21020208>.
- 446 Dore, C.M.P.G., Faustino Alves, M.G.d.C., Pofírio Will, L.S.E., Costa, T.G., Sabry, D.A.,
447 de Souza Rêgo, L.A.R., Accardo, C.M., Rocha, H.A.O., Filgueira, L.G.A., Leite, E.L.,
448 2013. A sulfated polysaccharide, fucans, isolated from brown algae *Sargassum vulgare*
449 with anticoagulant, antithrombotic, antioxidant and anti-inflammatory effects.
450 Carbohydr. Polym. 91, 467-475. <https://doi.org/10.1016/j.carbpol.2012.07.075>.
- 451 Fernández, P.A., Hurd, C.L., Roleda, M.Y., 2014. Bicarbonate uptake via an anion exchange
452 protein is the main mechanism of inorganic carbon acquisition by the giant kelp
453 *Macrocystis pyrifera* (Laminariales, Phaeophyceae) under variable pH. J. Phycol. 50,
454 998-1008. <https://doi.org/10.1111/jpy.12247>.
- 455 Fernández, P.A., Roleda, M.Y., Hurd, C.L., 2015. Effects of ocean acidification on the
456 photosynthetic performance, carbonic anhydrase activity and growth of the giant kelp
457 *Macrocystis pyrifera*. Photosynth. Res. 124, 293-304. [https://doi.org/10.1007/s11120-](https://doi.org/10.1007/s11120-015-0138-5)
458 015-0138-5.
- 459 Fernández, P.A., Roleda, M.Y., Rautenberger, R., Hurd, C.L., 2018. Carbonic anhydrase
460 activity in seaweeds: overview and recommendations for measuring activity with an
461 electrometric method, using *Macrocystis pyrifera* as model species. Mar. Biol. 165,
462 <https://doi.org/10.1007/s00227-018-3348-5>.
- 463 Figueroa, F. Malta, E. Bonomi-Barufi, J. Conde-Álvarez, R. Nitschke, U. Arenas, F., 2014.
464 Short-term effects of increasing CO₂, nitrate and temperature on three Mediterranean

- macroalgae: biochemical composition. *Aquat. Biol.* 22, 177-193. [https://doi: 10.3354/ab00610](https://doi.org/10.3354/ab00610).
- Foo, S.A., Byrne, M., Ricevuto, E., Gambi, M.C., 2018. The carbon dioxide vents of Ischia, Italy, a natural system to assess impacts of ocean acidification on marine ecosystems: an overview of research and comparisons with other vent systems. In: Hawkins, S.J., Evans, A.J., Dale, A.C., Flirth, L. B., Smith, I. P. (Eds.), *Oceanography and Marine Biology: An Annual Review*, 56, 237-310. <https://doi.org/10.1201/9780429454455>.
- Gao, G., Clare, A.S., Chatzidimitriou, E., Rose, C., Caldwell, G., 2018. Effects of ocean warming and acidification, combined with nutrient enrichment, on chemical composition and functional properties of *Ulva rigida*. *Food Chem.* 258, 71–78. <https://doi.org/10.1016/j.foodchem.2018.03.040>.
- Gao, G., Clare, A.S., Rose, C., Caldwell, G.S., 2017. Eutrophication and warming-driven green tides (*Ulva rigida*) are predicted to increase under future climate change scenarios. *Mar. Poll. Bull.* 144, 439-447. <https://doi.org/10.1016/j.marpolbul.2016.10.003>.
- Gao, G., Gao, Q., Bao, M., Xu, J., Li, X., 2019. Nitrogen availability modulates the effects of ocean acidification on biomass yield and food quality of a marine crop *Pyropia yezoensis*. *Food Chem.* 271, 623-629. <https://doi.org/10.1016/j.foodchem.2018.07.090>.
- Gao, K., Beardall, J., Häder, D.P., Hall-Spencer, J. M., Gao, G., Hutchins, D. A., 2019. Effects of ocean acidification on marine photosynthetic organisms under the concurrent influences of warming, UV radiation and deoxygenation. *Front. Mar. Sci.* 6, 322. <https://doi.org/10.3389/fmars.2019.00322>.
- Ghania, A., Nabila, B.B., Larbi, B., Elisabeth, M., Philippe, G., Mariem, B., Khadidja, K.K., Wacila, B.R., Fawzia, A.B., 2019. Antimicrobial and antiparasitic activities of three algae from the northwest coast of Algeria. *Nat. Prod. Res.* 33, 742-745. doi: 10.1080/14786419.2017.1405403.

- 491 Ghasemzadeh, A., Jaafar, Z.H., 2011. Antioxidant potential and anticancer activity of young
 492 ginger (*Zingiber officinale* Roscoe) grown under different CO₂ concentration. J. Med.
 493 Plants Res. 5, 3247-3255.
- 494 Graiff, A., Bartsch, I., Ruth, W., Wahl, M., Karsten, U., 2015. Season exerts differential
 495 effects of ocean acidification and warming on growth and carbon metabolism of the
 496 seaweed *fucusfocus vesiculosus* in the western Baltic Sea. Front. Microbiol. 2, 112.
 497 <https://doi.org/10.3389/fmars.2015.00112>.
- 498 Gutow, L., Rahman, M.M., Bartl, K., Saborowski, R., Bartsch, I., Wiencke, C., 2014. Ocean
 499 acidification affects growth but not nutritional quality of the seaweed *Fucus vesiculosus*
 500 (Phaeophyceae, Fucales). J. Exp. Mar. Biol. Ecol. 453, 84–90.
 501 <https://doi.org/10.1016/j.jembe.2014.01.005>.
- 502 Hallé, J.P., Landry, D., Fournier, A., Beaudry, M., Leblond, F.A., 1993. Method for the
 503 quantification of alginate in microcapsules. Cell Transplant. 2, 429-436.
 504 <https://doi.org/10.1177/096368979300200511>.
- 505 Hall-Spencer, J. M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner, S.M.,
 506 Rowley, S.J., Tedesco, D., Buia, M.C., Volcanic carbon dioxide vents show ecosystem
 507 effects of ocean acidification, Nature. 454, 96-99. <https://doi:10.1038/nature07051>.
- 508 Hao, H., Fu, M., Yan, R., He, B., Li, M., Liu, Q., Zhang, X., Huang, R., 2019. Chemical
 509 composition and immunostimulatory properties of green alga *Caulerpa racemosa* var
 510 peltata. Food Agr. Immunol. 30, 937-954.
 511 <https://doi.org/10.1080/09540105.2019.1646216>.
- 512 Hayatsu, H., Negishi, T., Arimoto, S., Hayatsu, T., 1993. Porphyrins as potential inhibitors
 513 against exposure to carcinogens and mutagens. Mutat Res-Fund Mol. M. 290, 79-85.
 514 [https://doi.org/10.1016/0027-5107\(93\)90035-E](https://doi.org/10.1016/0027-5107(93)90035-E).

- 515 Hofmann, L.C., Straub, S., Bischof, K., 2013. Elevated CO₂ levels affect the activity of
 516 nitrate reductase and carbonic anhydrase in the calcifying rhodophyte *Corallina*
 517 *officinalis*. J. Exp. Bot. 64 (4), 899–908. <https://doi.org/10.1093/jxb/ers369>.
- 518 Idso, S.B., Kimball, B.A., Pettit III, G.R., Garner, L.C., Pettit, G.R., Backhaus, R.A., 2000.
 519 Effects of atmospheric CO₂ enrichment on the growth and development of *Hymeno*
 520 *callislittoralis* (Amaryllidaceae) and the concentrations of several antineoplastic and
 521 antiviral constituents of its bulbs. Am. J. Bot. 87, 769-773.
 522 <https://doi.org/10.2307/2656884>.
- 523 Jaafar, H.Z.E., Ibrahim, M.H., Karimi, E., 2012. Phenolics and Flavonoids Compounds,
 524 Phenylalanine Ammonia Lyase and Antioxidant Activity Responses to Elevated CO₂ in
 525 *Labisia pumila* (Myrsinaceae). Molecules 17, 6331.
 526 <https://doi.org/10.3390/molecules17066331>.
- 527 Jun, J.Y., Jung, M.J., Jeong, I.H., Yamazaki, K., Kawai, Y., Kim, B.M., 2018. Antimicrobial
 528 and antibiofilm activities of sulfated polysaccharides from marine algae against dental
 529 plaque bacteria. Mar. Drugs. 16, 301. <https://doi.org/10.3390/md16090301>.
- 530 Koch, M., Bowes, G., Ross, C., Zhang, X.H., 2013. Climate change and ocean acidification
 531 effects on seagrasses and marine macroalgae. Glob. Change Biol. 19, 103-132.
 532 <https://doi.org/10.1111/j.1365-2486.2012.02791.x>.
- 533 Kolsi, R.B.A., Salah, H.B., Jardak, N., Chaaben, R., Jribi, I., Feki, A.E., Rebai, T., Jamoussi,
 534 K., Allouche, N., Blecker, C., Belghith, H., Belghith, K., 2017. Sulphated
 535 polysaccharide isolated from *Sargassum vulgare*: characterization and hypolipidemic
 536 effects. Carbohydr. Polym. 170, 148-159. <https://doi.org/10.1016/j.carbpol.2017.04.083>.
- 537 Kroeker, K.J., Micheli, F., Gambi, M.C., 2013. Ocean acidification causes ecosystem shifts
 538 via altered competitive interactions. Nat. Clim. Change. 3, 156-159.
 539 <https://doi.org/10.1038/nclimate1680>.

- 540 Kübler, J.E., Johnston, A.M., Raven, J.A., 1999. The effects of reduced and elevated CO₂
 541 and O₂ on the seaweed *Lomentaria articulata*. Plant Cell Environ. 22, 1303-1310.
 542 <https://doi.org/10.1046/j.1365-3040.1999.00492.x>.
- 543 Kumar, A., Abdelgawad, H., Castellano, I., Lorenti, M., Delledonne, M., Beemster, G.T.S.,
 544 Asard, H., Buia, M.C., Palumbo, A., 2017a. Physiological and biochemical analyses
 545 shed light on the response of *Sargassum vulgare* to ocean acidification at different time
 546 scales. Front. Plant Sci. 8, 570-570. <https://doi.org/10.3389/fpls.2017.00570>.
- 547 Kumar, A., Abdelgawad, H., Castellano, I., Selim, S., Beemster, G.T.S., Asard, H., Buia,
 548 M.C., Palumbo, A., 2018. Effects of ocean acidification on the levels of primary and
 549 secondary metabolites in the brown macroalga *Sargassum vulgare* at different time
 550 scales. Sci. Total Environ. 643, 946-956.
 551 <https://doi.org/10.1016/j.scitotenv.2018.06.176>.
- 552 Kumar, A., Castellano, I., Patti, F.P., Delledonne, M., Abdelgawad, H., Beemster, G.T.S.,
 553 Asard, H., Palumbo, A., Buia, M.C., 2017b. Molecular response of *Sargassum vulgare*
 554 to acidification at volcanic CO₂ vents: insights from *de novo* transcriptomic analysis.
 555 Mol. Ecol. 26, 2276-2290. <https://doi.org/10.1111/mec.14034>.
- 556 Leakey, A.D.B., Xu, F., Gillespie, K.M., McGrath, J.M., Ainsworth, E.A., Ort, D.R., 2009.
 557 Genomic basis for stimulated respiration by plants growing under elevated carbon
 558 dioxide. Proc. Natl. Acad. Sci., 106, 3597-3602.
 559 <https://doi.org/10.1073/pnas.0810955106>.
- 560 Liu, L., Heinrich, M., Myers, S., Dworjanyn, S.A., 2012. Towards a better understanding of
 561 medicinal uses of the brown seaweed *Sargassum* in traditional Chinese medicine: A
 562 phytochemical and pharmacological review. J. Ethnopharmacol. 142, 591-619.
 563 <https://doi.org/10.1016/j.jep.2012.05.046>.
- 564 Lombardi, C., Rodolfo-Metalpa, R., Cocito, S., Gambi, M.C., Taylor, P.D., 2011. Seawater
 565 carbonate chemistry and MgCO₃ concentration in bryozoan *Myriapora truncata*

- 566 branches during experiments. Mar. Ecol. 32, 211-22.
 567 <https://doi.org/10.1594/PANGAEA.771573>.
- 568 Lorenti, M., Buia, M.C., Di Martino, V., Modigh, M., 2005. Occurrence of mucous
 569 aggregates and their impact on *Posidonia oceanica* beds. Sci. Total Environ. 353, 369–
 570 379. <https://doi.org/10.1016/j.scitotenv.2005.09.025>.
- 571 Lorenti, M., De Falco, G., 2004. Measurements and characterization of abiotic variables. In:
 572 Gambi, M.C., Dappiano, M. (Eds), Mediterranean Marine Benthos: A Manual of
 573 Methods for its Sampling and Study pp. 1–37.
- 574 Machu, L., Misurcova, L., VavraAmbrozova, J., Orsavova, J., Mlcek, J., Sochor, J.,
 575 Jurikova, T., 2015. Phenolic content and antioxidant capacity in algal food products.
 576 Molecules 20, 1118-33. <https://doi.org/10.3390/molecules20011118>.
- 577 Mackey, K.R.M., Morris, J.J., Morel, F.M.M., Kranz, S.A., 2015. Response of
 578 photosynthesis to ocean acidification. Oceanography 28, 74-91.
 579 <https://doi.org/10.5670/oceanog.2015.33>.
- 580 Maron, D.M., Ames, B.N., 1983. Revised methods for the Salmonella mutagenicity test.
 581 Mutat. Res-Envir. Muta. 113, 173-215. [tps://doi.org/10.1016/0165-1161\(83\)90010-9](https://doi.org/10.1016/0165-1161(83)90010-9).
- 582 Mouritsen, O.G., Rhatigan, P., Pérez-Lloréns, J.L., 2018. The rise of seaweed gastronomy:
 583 phycogastronomy. Bot. Mar. 62, 195-209. <https://doi.org/10.1515/bot-2018-0041>.
- 584 Ohkawa, H., Ohishi, N., Yagi, K., 1979. Assay for lipid peroxides in animal tissues by
 585 thiobarbituric acid reaction. Anal. Biochem. 95, 351–358. [https://doi.org/10.1016/0003-](https://doi.org/10.1016/0003-2697(79)90738-3)
 586 2697(79)90738-3.
- 587 Osuna-Ruiz, I., López-Saiz, C.M., Burgos-Hernández, A., Velázquez, C., Nieves-Soto, M.,
 588 Hurtado-Oliva, M. A., 2016. Antioxidant, antimutagenic and antiproliferative activities
 589 in selected seaweed species from Sinaloa, Mexico. Pharm. Biol. 54, 2196-2210.
 590 <https://doi.org/10.3109/13880209.2016.1150305>.

- 591 Paiva, L., Lima, E., Neto, A., Baptista, J., 2018. Seasonal variability of the biochemical
 592 composition and antioxidant properties of *Fucus spiralis* at two Azorean Islands. Mar.
 593 Drugs 16, 248. <https://doi.org/10.3390/md16080248>.
- 594 Perumal, B., Chitra, R., Maruthupandian, A., Viji, M., 2019. Nutritional assessment and
 595 bioactive potential of *Sargassum polycystum* C. Agardh (Brown Seaweed), Indian J
 596 Geo. Mar. Sci. 48, 492-498.
- 597 Porzio, L., Buia, M.C., Hall-Spencer, J.M., 2011. Effects of ocean acidification on
 598 macroalgal communities. J. Exp. Mar. Biol. Ecol. 400, 278-287.
 599 <https://doi.org/10.1016/j.jembe.2011.02.011>.
- 600 Porzio, L., Buia, M. C., Lorenti, M., De Maio, A., Arena, C., 2017. Physiological responses
 601 of a population of *Sargassum vulgare* (Phaeophyceae) to high pCO₂/low pH:
 602 implications for its long-term distribution. Sci. Total Environ. 576, 917-925.
 603 <https://doi.org/10.1016/j.scitotenv.2016.10.096>.
- 604 Praiboon, J., Palakas, S., Noiraksa, T., Miyashita, K., 2018. Seasonal variation in nutritional
 605 composition and anti-proliferative activity of brown seaweed, *Sargassum oligocystum*.
 606 J. Appl. Phycol. 30, 101-111. <https://doi.org/10.1007/s10811-017-1248-6>.
- 607 Pramesti, R., Setyati, W.A., Zainuddin, M., Puspita, M., 2017. Bioecology of *Sargassum* sp.
 608 and its extract bioactivity as Anti-MDR Bacteria. Ilmu Kelautan. 22, 8.
 609 <https://doi.org/10.14710/ik.ijms.22.4.185-192>.
- 610 Ráz, B., Iten, M., Grether-Bühler, Y., Kaminsky, R., Brun, R., 1997. The Alamar Blue®
 611 assay to determine drug sensitivity of *African trypanosomes* (*T.b. rhodesiense* and *T.b.*
 612 *gambiense*) in vitro. Acta Trop. 68, 139-147. [https://doi.org/10.1016/S0001-](https://doi.org/10.1016/S0001-706X(97)00079-X)
 613 [706X\(97\)00079-X](https://doi.org/10.1016/S0001-706X(97)00079-X).
- 614 d'Alcalà, M.R., Conversano, F., Corato, F., Licandro, P., Mangoni, O., Marino, D.,
 615 Mazzocchi, M.G., Modigh, M., Montresor, M., Nardella, M. and Saggiomo, V., 2004.
 616 Seasonal patterns in plankton communities in a pluriannual time series at a coastal

- 617 Mediterranean site (Gulf of Naples): an attempt to discern recurrences and trends. Sci.
 618 Mar. 68, 65–83. <https://doi.org/10.3989/scimar.2004.68s165>.
- 619 Ricevuto, E., Kroeker, K., Ferrigno, F., Micheli, F., Gambi, M., 2014. Spatio-temporal
 620 variability of polychaete colonization at volcanic CO₂ vents indicates high tolerance to
 621 ocean acidification. Mar. Biol. 161, 2909-2919. [https://doi.org/10.1007/s00227-014-](https://doi.org/10.1007/s00227-014-2555-y)
 622 2555-y.
- 623 Ricevuto, E., Salvatrice V, Maria.Cristina G. 2015. Ocean acidification effects on stable
 624 isotope signatures and trophic interactions of polychaete consumers and organic matter
 625 sources at a CO₂ shallow vent system. J. Exp. Mar. Biol. Ecol. 468, 105-117.
 626 <https://doi.org/10.1016/j.jembe.2015.03.016>.
- 627 Roleda M.Y., Hurd C.L., 2012. Seaweed Responses to Ocean Acidification, In: Wiencke C.,
 628 Bischof K. (Eds.), Seaweed Biology: Ecological Studies (Analysis and Synthesis).
 629 Springer, Berlin, pp. 407-432. <https://doi.org/10.1007/978-3-642-28451-9>.
- 630 Roleda, M.Y., Marfaing, H., Desnica, N., Jónsdóttir, R., Skjermo, J., Rebours, C., Nitschke,
 631 U., 2019. Variations in polyphenol and heavy metal contents of wild-harvested and
 632 cultivated seaweed bulk biomass: Health risk assessment and implication for food
 633 applications. Food Control 95,121-134. <https://doi.org/10.1016/j.foodcont.2018.07.031>.
- 634 Roleda, M.Y., Morris, J.N., McGraw, C.M., Hurd, C.L., 2012. Ocean acidification and
 635 seaweed reproduction: increased CO₂ ameliorates the negative effect of lowered pH on
 636 meiospore germination in the giant kelp *Macrocystis pyrifera* (Laminariales,
 637 Phaeophyceae). Glob. Chang. Biol. 18, 854–864. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2486.2011.02594.x)
 638 2486.2011.02594.x.
- 639 Saleh, A.M., Selim, S., Jaouni, S.A., AbdElgawad, H., 2018. CO₂ enrichment can enhance
 640 the nutritional and health benefits of parsley (*Petroselinum crispum* L.) and dill
 641 (*Anethum graveolens* L.). Food Chem. 269, 519-526.
 642 <https://doi.org/10.1016/j.foodchem.2018.07.046>.

- 643 Selim, S.A., Mohamed, H.A.A., Mona, S.M., Warrad, M.F., 2013. Antibacterial activities,
 644 chemical constituents and acute toxicity of Egyptian *Origanum majorana* L., *Peganum*
 645 *harmala* L.L. and *Salvia officinalis* L. essential oils. Afr. J. Pharm. Pharmacol. 7, 725-
 646 735. <https://doi.org/10.5897/AJPP2013.3518>.
- 647 Serebryakova, A., Aires, T., Viard, F., Serrão, E.A., Engelen, A.H., 2018. Summer shifts of
 648 bacterial communities associated with the invasive brown seaweed *Sargassum muticum*
 649 are location and tissue dependent. Plos One 13, e0206734.
 650 <https://doi.org/10.1371/journal.pone.0206734>.
- 651 Singh, R.P., Reddy, C.R.K., 2014. Seaweed–microbial interactions: key functions of
 652 seaweed-associated bacteria. FEMS Microbiol. Ecol. 88, 213-230.
 653 <https://doi.org/10.1111/1574-6941.12297>.
- 654 Smit, A.J., 2004. Medicinal and pharmaceutical uses of seaweed natural products: a review.
 655 J Appl. Phycol. 16, 245-262. <https://doi.org/10.1023/B:JAPH.0000047783.36600.ef>.
- 656 Solowey, E., Lichtenstein, M., Sallon, S., Paavilainen, H., Solowey, E., Lorberboum-Galski,
 657 H., 2014. Evaluating Medicinal Plants for Anticancer Activity. Scientific World
 658 Journal. Article id: 721402. <http://dx.doi.org/10.1155/2014/721402>.
- 659 Sousa, A.d.P.A., Barbosa, P.S.F., Torres, M.R., Martins, A.M.C., Martins, R.D., Alves,
 660 R.d.S., Sousa, D.F.d., Alves, C.D., Costa-Lotufo, L.V., Monteiro, H.S.A., 2008. The
 661 renal effects of alginates isolated from brown seaweed *Sargassum vulgare*. J. Appl.
 662 Toxicol. 28, 364-369. <https://doi.org/10.1002/jat.1287>.
- 663 Suárez-Álvarez, S., Gómez-Pinchetti, J.L., García-Reina, G., 2012. Effects of increased CO₂
 664 levels on growth, photosynthesis, ammonium uptake and cell composition in the
 665 macroalga *Hypnea spinella* (Gigartinales, Rhodophyta). J. Appl. Phycol. 24, 815-823.
 666 <https://doi.org/10.1007/s10811-011-9700-5>.

- 667 Torres, F.A.E., Passalacqua, T.G., Velásquez, A.M.A., de Souza, R.A., Colepicolo, P.,
 668 Graminha, M.A.S., 2014. New drugs with antiprotozoal activity from marine algae: a
 669 review. *Rev. Bras. Farmacogn.* 24, 265-276. <https://doi.org/10.1016/j.bjp.2014.07.001>.
- 670 Torres, M.R., Sousa, A.P.A., Silva Filho, E.A.T., Melo, D.F., Feitosa, J.P.A., de Paula,
 671 R.C.M., Lima, M.G.S., 2007. Extraction and physicochemical characterization of
 672 *Sargassum vulgare* alginate from Brazil. *Carbohydr. Res.* 342, 2067-2074.
 673 <https://doi.org/10.1016/j.carres.2007.05.022>.
- 674 Turan, G., Cirik, S., Tekogul, H., Koru, E., Karacalar, U., Seyhaneyildiz, S., 2017.
 675 Determination of the Seasonal Yields of Total Fucose and Fucoidan Yields in Brown
 676 Sea- weeds (Order Fucales) Distributed along the Coast of Urla (Izmir, Turkey). *J*
 677 *Aquac. Fisheries.* 1, 005. <https://doi: 10.24966/AAF-5523/100005>
- 678 Véliz, K., Chandía, N., Karsten, U., Lara, C., Thiel, M., 2019. Geographic variation in
 679 biochemical and physiological traits of the red seaweeds *Chondracanthus chamissoi*
 680 and *Gelidium lingulatum* from the south east Pacific coast. *J. Appl. Phycol.* 31, 665-
 681 682. <https://doi.org/10.1007/s10811-018-1532-0>.
- 682 Wu, H., Zou, D., Gao, K., 2008. Impacts of increased atmospheric CO₂ concentration on
 683 photosynthesis and growth of micro- and macro-algae. *Sci. China Ser. C Life Sci.*
 684 51,1144–1150. <https://doi.org/10.1007/s11427-008-0142-5>.
- 685 Wu, L., Sun, J., Su, X., Yu, Q., Yu, Q., Zhang, P., 2016. A review about the development of
 686 fucoidan in antitumor activity: Progress and challenges. *Carbohydr. Polym.* 154, 96-
 687 111. <https://doi.org/10.1016/j.carbpol.2016.08.005>.
- 688 Yousefzadi, M., Mirjalili, M.H., 2017. Effect of Climate Change on Algae Valuable Source
 689 of Medicinal Natural Compounds. In: Ghorbanpour, M., Varma, A. (Eds.), *Medicinal*
 690 *Plants and Environmental Challenges.* Springer, USA pp. 123-135.
 691 https://doi.org/10.1007/978-3-319-68717-9_7.
- 692

Figure Legends

Fig. 1. Study sites around the coast of the Ischia Island, Italy: Castello Aragonese (CO₂ vents; acidified site) and Lacco Ameno (control site; Image generated through QGIS version 2.12.2).

Fig. 2. Concentration of Fucoidan and Alginate in the extract of *S. vulgare* collected at the control and the acidified site. Values are mean $\pm$ SE, n= 3, ***p <0.001.

Fig. 3. *In vitro* antioxidant activity of the extract of *S. vulgare* collected at the control and the acidified site, A: antioxidant capacity using FRAP (ferric reducing/antioxidant power) assay; B. antioxidant activity using DPPH (diphenylpicrylhydrazyl) radical scavenging assay; C. anti-lipid peroxidation assay. Values are mean $\pm$ SE, n=3, *** p < 0.001.

Fig. 4. Antimicrobial activity of *S. vulgare* collected from the control and acidified site against human pathogens. Values are mean $\pm$ SE, n=3, *p< 0.05, ** p< 0.01, ***p< 0.001.

Fig. 5. Antiprotozoal activity of *S. vulgare* collected from the control and acidified site against *T. acruzi* and *L. donovani*. Values are mean $\pm$ SE, n=3, *p< 0.05, ** p< 0.01, ***p< 0.001.

Fig. 6. Anticancer activity of *S. vulgare* collected from the control and acidified site against four cancer cell lines hepatocellular carcinoma (HepG2), colon carcinoma (Colo205), embryonic kidney adenocarcinoma (293), and urinary bladder carcinoma (T24P). Values are mean $\pm$ SE, n=3, *p< 0.05, ** p< 0.01, ***p< 0.001.

Fig. 7. Antimutagenic activity of *S. vulgare* collected from control and acidified site against *S. typhimuriam* TA 98 and TA 100. Values are mean $\pm$ SE, n=3, *p< 0.05, ** p< 0.01, ***p< 0.001.

Appendices Table A. 1: Summary of water chemistry at the control site along Lacco Ameno and at the acidified site along Castelo Aragonese, Ischia Island, where *S. Vulgare* occurs.

¹ Hall-Spencer *et al.*, 2008; ² Cigliano *et al.*, 2010; ³ Frieder *et al.*, 2013; ⁴ Kroeker *et al.*, 2011; ⁵ Calosi *et al.*, 2013; ⁶ Ricevuto *et al.*, 2014. *Point value on 29th sept 2014, obtained from Dr. Gil Rilov.

	Acidified site, Castello Aragonese							Control stie, Lacco Ameno
Parameters	2007 (Apr-May) ¹	2008 (Apr-May) ²	2009 (Sep) ³	2010(Sep-Oct) ⁴	2011 (May-June) ⁵	2013 (Apr-May) ⁶	2014 (Aug-Sep)	2013 (Apr-May) ⁶
Salinity	38	38	38.1±0.04	37±0.2	37.14±0.05	37.11±0.6	37±0.5	36.9±0.7
Temperature	18.5	19.44±1.98	25±0.63	17.5±2.8	23.31±0.12	19.02±1.21	15±0.4	19.10±1.23
pH	6.57±0.06	7.32±0.29	6.83±0.42	6.59±0.51	7.07±0.07	7.14±0.4	6.80±0.09	8.13±0.05
TA(µequi/kg)	2500	2566±12.98	2578±6.1	2563±13	2673.73±12.57	2597.40±75	2608*	2612.93±26.5
pCO ₂ (µatm)	20812±2253	3317.80±1744	15099±2997	23989±16638	5337.65±567.54	4266.2±2814	NA	165.8±24.6
HCO ₃ ⁻ (µmol/kg)	2470±100	2418±113	NA	NA	2565.58±20.72	2518.9±101.4	NA	2067.3±53.9
CO ₃ ²⁻ (µmol/kg)	100±10	62±44.94	NA	NA	38.30±2.93	31±40.1	NA	217.2±21.8
DIC(µmol/kg)	3160±80	2585.20±122.68	2564±108	3849±790	2781.21±20.15	2549.92±61.30	NA	2284.55±32.09

Appendices Table A2. T- test statistics summary for bioactivity of *S. vulgare* from acidified and control site.


Independent Samples Test										
		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Antiprotozoal_TA	Equal variances assumed	0.698	0.435	5.737	6.000	0.001218	2.627	0.458	1.507	3.747
	Equal variances not assumed			5.737	5.166	0.002023	2.627	0.458	1.461	3.793
Antiprotozoal_LD	Equal variances assumed	1.234	0.309	3.007	6.000	0.002379	1.460	0.485	0.272	2.647
	Equal variances not assumed			3.007	5.163	0.028679	1.460	0.485	0.224	2.695
Antibac_Strepto	Equal variances assumed	1.797	0.229	-2.926	6.000	0.002643	-2.164	0.740	-3.974	-0.354
	Equal variances not assumed			-2.926	3.966	0.043440	-2.164	0.740	-4.224	-0.104


Antibac_E.coli	Equal variances assumed	0.636	0.456	- 4.670	6.000	0.003432	-5.774	1.236	-8.799	-2.748
	Equal variances not assumed			- 4.670	5.034	0.005392	-5.774	1.236	-8.945	-2.602
Antibac_Serretia	Equal variances assumed	2.029	0.204	- 2.514	6.000	0.045649	-2.991	1.190	-5.902	-0.080
	Equal variances not assumed			- 2.514	3.542	0.073804	-2.991	1.190	-6.469	0.487
Antibac_S.aureus	Equal variances assumed	0.029	0.871	- 2.402	6.000	0.053133	-4.043	1.683	-8.162	0.075
	Equal variances not assumed			- 2.402	5.778	0.054716	-4.043	1.683	-8.200	0.114
Antibac_K.pneumoniae	Equal variances assumed	0.153	0.709	- 3.862	6.000	0.008338	-4.955	1.283	-8.094	-1.816
	Equal variances not assumed			- 3.862	5.421	0.010143	-4.955	1.283	-8.177	-1.733
Antibac_P.vulgaris	Equal variances assumed	0.500	0.506	- 5.044	6.000	0.002347	-6.211	1.231	-9.223	-3.198
	Equal variances not			- 5.044	4.708	0.004671	-6.211	1.231	-9.436	-2.986


	assumed									
Antibac_P.aeruginosa	Equal variances assumed	0.001	0.976	-8.662	6.000	0.000131	-4.268	0.493	-5.474	-3.062
	Equal variances not assumed			-8.662	5.998	0.000131	-4.268	0.493	-5.474	-3.062
Antifungal_C.albicans	Equal variances assumed	0.947	0.368	-0.665	6.000	0.530621	-1.123	1.688	-5.254	3.008
	Equal variances not assumed			-0.665	4.316	0.539770	-1.123	1.688	-5.678	3.432
Antifungal_A.parasiticus	Equal variances assumed	1.305	0.297	-7.341	6.000	0.000327	-4.556	0.621	-6.075	-3.037
	Equal variances not assumed			-7.341	3.610	0.002708	-4.556	0.621	-6.355	-2.757
Antifungal_P.expansum	Equal variances assumed	1.437	0.276	-2.164	6.000	0.073681	-4.606	2.128	-9.814	0.602
	Equal variances not assumed			-2.164	4.144	0.094086	-4.606	2.128	-10.435	1.224
Antifungal_T.harzia num	Equal variances assumed	0.088	0.777	-0.618	6.000	0.559508	-0.238	0.385	-1.181	0.705


	Equal variances not assumed			- 0.618	5.311	0.562335	-0.238	0.385	-1.211	0.735
Antifungal_F.oxysporum	Equal variances assumed	1.661	0.245	- 3.812	6.000	0.008845	-3.969	1.041	-6.516	-1.421
	Equal variances not assumed			- 3.812	5.316	0.011116	-3.969	1.041	-6.598	-1.340
Antimutagenic_S.typhimurium TA98	Equal variances assumed	1.543	0.261	- 5.354	6.000	0.001738	-11.767	2.198	-17.144	-6.389
	Equal variances not assumed			- 5.354	3.664	0.007500	-11.767	2.198	-18.096	-5.437
Antimutagenic_S.typhimuriumTA100	Equal variances assumed	0.494	0.509	- 2.754	6.000	0.033137	-8.626	3.133	-16.291	-0.961
	Equal variances not assumed			- 2.754	5.212	0.038401	-8.626	3.133	-16.581	-0.671
AntiCancer_HepG2	Equal variances assumed	1.981	0.209	- 8.483	6.000	0.000147	-25.583	3.016	-32.962	-18.203
	Equal variances not assumed			- 8.483	3.432	0.002028	-25.583	3.016	-34.532	-16.633
Anticancer_Colo205	Equal variances	2.660	0.154	- 8.438	6.000	0.000151	-13.828	1.639	-17.838	-9.818


	assumed									
	Equal variances not assumed			- 8.438	3.445	0.002032	-13.828	1.639	-18.682	-8.974
Anticance_293	Equal variances assumed	0.534	0.492	- 5.086	6.000	0.000225	-4.307	0.847	-6.379	-2.235
	Equal variances not assumed			- 5.086	5.063	0.003683	-4.307	0.847	-6.475	-2.138
Anticancer_T24P	Equal variances assumed	1.907	0.216	- 5.194	6.000	0.000202	-10.249	1.973	-15.078	-5.421
	Equal variances not assumed			- 5.194	3.439	0.009798	-10.249	1.973	-16.099	-4.399

















Highlights

- Bioactivities of *S. vulgare* from Ischia CO₂ vents and nearby control site were analysed.
- Elevated DIC increases polysaccharide content in the algae at CO₂ vents.
- Algal extract from acidified population showed higher antimicrobial, and antiprotozoal activity.
- Acidified population showed pronounced antimutagenic potential and anticancer activities.

AUTHORSHIP STATEMENT

Manuscript title: **Ocean acidification affects biological activities of seaweeds: A case study of *Sargassum vulgare* from Ischia volcanic CO₂ vents**

All persons who meet authorship criteria are listed as authors, and all authors certify that they have participated sufficiently in the work to take public responsibility for the content, including participation in the concept, design, analysis, writing, or revision of the manuscript. Furthermore, each author certifies that this material or similar material has not been and will not be submitted to or published in any other publication before its appearance in the *Hong Kong Journal of Occupational Therapy*.

Authorship contributions

Please indicate the specific contributions made by each author (list the authors' initials followed by their surnames, e.g., Y.L. Cheung). The name of each author must appear at least once in each of the three categories below.

Category 1

Conception and design of study: **A. Kumar, M.C. Buia, A. Palumbo, H. AbdElgawad**

acquisition of data: **M. Mohany, M.A.M. Wadaan, W.N. Hozzein, H. AbdElgawad**

analysis and/or interpretation of data: **A. Kumar, H. AbdElgawad, G.T.S. Beemster**

Category 2

Drafting the manuscript: **A. Kumar**

revising the manuscript critically for important intellectual content: **M.C. Buia, A. Palumbo, H. AbdElgawad, G.T.S. Beemster**

Category 3

Approval of the version of the manuscript to be published (the names of all authors must be listed):


Amit Kumar, Maria Cristina Buia, Anna Palumbo, Mohamed Mohany, Mohammed A.M.

Wadaan, Wael N. Hozzein, Gerrit T.S. Beemster, Hamada AbdElgawad

Acknowledgements

All persons who have made substantial contributions to the work reported in the manuscript (e.g., technical help, writing and editing assistance, general support), but who do not meet the criteria for authorship, are named in the Acknowledgements and have given us their written permission to be named. If we have not included an Acknowledgements, then that indicates that we have not received substantial contributions from non-authors.

This statement is signed by all the authors (a photocopy of this form may be used if there are more than 10 authors):

Author's name (typed)	Author's signature	Date
Amit Kumar		15/11/2019
Maria Cristina Buia		15/11/2019
Anna Palumbo		15/11/2019
Mohamed Mohany		15/11/2019
Mohammed A.M. Wadaan		15/11/2019
Wael N. Hozzein		15/11/2019
Gerrit T.S. Beemster		15/11/2019
Hamada AbdElgawad		15/11/2019

Declaration of interests

- ☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.
- ☒ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Amit Kumar
Maria Cristina Buia
Anna Palumbo
Mohamed Mohany
Mohammed A.M. Wadaan
Wael N. Hozzein
Hamada AbdElgawad